

JOURNAL OF AGRICULTURAL RESEARCH

Vol. XV

WASHINGTON, D. C., DECEMBER 23, 1918

No. 12

PARASITISM, MORPHOLOGY, AND CYTOLOGY OF *CRONARTIUM RIBICOLA*

By REGINALD H. COLLEY

Assistant Pathologist, Investigations in Forest Pathology, United States Department of
Agriculture

INTRODUCTION

During the last few years *Cronartium ribicola* Fischer has become the most serious fungus pest of white pine (*Pinus strobus* L.) in America, and as such it has been the subject of much study by pathologists who realized, from a knowledge of the course of the parasite in Europe, the damage it was capable of causing. Spaulding (54)¹ in the most comprehensive paper on the fungus which has appeared in this country, reviews the reports of its ravages on white pine in the Old World and gives its general characters and life history. Since the discovery by Stewart (57) of *C. ribicola* on species of *Ribes* at Geneva, N. Y., in 1906, many papers have appeared calling attention to the absolute necessity of controlling the spread of this parasite if the white pine is to be saved for reforestation. A great deal of attention has been given to testing the susceptibility of possible hosts through inoculation and to control and eradication methods, both in the United States and Canada. In Europe most of the recent work has been along this same line. Klebahn (23, 24, 25, 26) and Tubeuf (58, 59, 60) have carried on extensive experiments.

The morphology of *Cronartium ribicola* and the details of the interrelations of the parasite and its hosts have never been thoroughly worked out. The cytology of the genus *Cronartium* has up to the present time received very little attention. In the following paper the results of certain observations extending over a period of two years will be presented, first, with reference to the minute histology of the fungus and the interrelations of host and parasite, and second, with reference to the cytological phenomena accompanying spore production in the different types of sori. The paper is offered as a contribution to our knowledge of the parasitism, morphology, and cytology of the genus *Cronartium*.

¹ Reference is made by number (italic) to "Literature cited," p. 655-659.

LIFE HISTORY AND HOSTS

The life history of *Cronartium ribicola* has been so well reported by Tubeuf (59, 60), Klebahn (25), Spaulding (54, 55), and others that it is only necessary to repeat it in outline by way of introduction to what follows. Sporidia from the teliospores produced on *Ribes* spp. infect young stems and branches of *Pinus strobus* and other 5-needled pines. The pycnial and æcial stages subsequently develop on the pines. Æcial spores from the pine infect leaves of *Ribes* spp., on which the uredinia are shortly formed. These sori are produced in successive generations throughout the summer. Telia develop from old uredinia, or as separate entities, in the form of compact columns. The teliospores germinate *in situ*, each one producing a promycelium which gives rise to four sporidia. A diagram of the life cycle is presented in text figure 1.¹

In the United States and Canada the pine most frequently attacked is *Pinus strobus*, although *P. flexilis* James and *P. parviflora* Sieb. and Zucc. have been found infected. Practically every known species of *Ribes* is susceptible to infection to a greater or less degree, and therefore the discovery of an immune variety is much to be desired. The results of inoculations on *Ribes* spp. in America have been reported by Spaulding and Gravatt (56), and further work is being conducted with all the species and varieties of *Ribes* obtainable.

EXPERIMENTAL METHODS

The method used for rapid examination of specimens of pine suspected of being infected has been described in detail in a previous paper (7). This procedure in brief is as follows: Sections from fresh pine bark are cut on an ether freezing microtome, rinsed in water, stained in safranin and *Lichtgrün*, cleared in clove oil followed by xylol, and mounted in balsam. They are then examined, preferably with an oil-immersion lens, to determine the presence or absence of the characteristic mycelium and the striking haustoria of the rust, the latter being especially important from a diagnostic standpoint. This method yields transparent sections which for general morphological study have not been surpassed by following any of the more complicated methods given below. Very little shrinkage occurs in the mycelium, and as the method is usually employed before pycnia and æcia appear, the preservation of the hyphæ in as near their natural shape as possible is practically all that is necessary. Moreover the host tissue shrinks so slightly that the distortion is entirely negligible.

Two killing agents were used in the preparation of material for paraffin or celloidin embedding: (1) formalin-alcohol, made by adding 6 cc. of full strength commercial formalin (U. S. P. VIII) to 94 cc. of 70 per cent

¹ The drawings for the text figures were made by the aid of a camera lucida and a special projection apparatus for the drawings on the plates a camera lucida was used. The photographs and photomicrographs

alcohol; and (2) Flemming's fluid, both weak and strong. The first was used for pine material only; the second for both pine and *Ribes* spp. Formalin-alcohol gave excellent result not only for gross mycelial characters but also in some cases for nuclear phenomena. The material sec-




FIG. 1.—Diagram representing the life cycle of *Cronartium ribicola*. The spores are drawn to approximately the same scale and are numbered in the order of their appearance: 1, pycniospore; 2, aeciospore; 3, urediniospore; 4, teliospore; 5, sporidium. The repetition of the figure 3 in the broken line between the urediniospore and the teliospore indicates that the urediniospores appear in successive generations in a single growing season. a, A young pine branch at the time infection usually takes place; b, pycnia; c, aecia; d, teliospores; e, uredinia; f, the dots on the currant leaf represent uredinia; g, the dots represent uredinia and the dashes telial columns; h, this figure represents the condition of the infected leaf in the fall when the telial columns are the predominant spore forms.

tioned easily, either in paraffin or celloidin, and stained well with safranin and *Lichtgrün* or Haidenhain's iron-alum hematoxylin. Both concentrations of Flemming's fluid were very satisfactory. The material was

thoroughly washed in running water. Certain generally accepted principles of killing and fixing were considerably modified by greatly shortening the time the material was left in the different reagents. The time the material remained in the grade alcohols was cut to 5 minutes for 10 per cent alcohol, 10 minutes for 15 per cent, 20 minutes for 20 per cent, and 30 minutes each for 30, 40, 50, 70, 80, 90, and 95 per cent. Absolute alcohol was used for 30 minutes with one change.

Xylol-alcohol was used both in short steps of 5, 10, 15, 20, 25, 50, and 75 per cent, or in a 25, 50, and 75 per cent concentration. For the short jumps the schedule of 5 minutes for 5 per cent, 10 minutes for 10 per cent, etc., was adopted; for all the longer jumps 30 minutes were allowed. The longer jumps apparently did not injure the material or its staining qualities. Pure xylol was used for one hour, with one change. The infiltration with paraffin was carried out as rapidly as was feasible, starting with shavings of 45° paraffin in xylol in cold solution, for 12 to 15 hours (overnight), then at the water-bath temperature, gradually adding more soft paraffin, for 24 hours. The mixture of xylol and paraffin was replaced by pure 45° paraffin for two hours, the latter by 55° paraffin for four hours, with one change and the material then embedded in the harder paraffin.

Æciospores and telial columns with promycelia and sporidia attached were killed in Flemming's fluid and handled with the aid of a centrifuge. By using a short spinning at moderate speed to send the small objects to the bottom of the centrifuge tube little material was lost in decanting or pipetting off the different reagents. When the material had been brought to pure paraffin, it was transferred to a glass tube 2 inches long with ¼-inch bore, which had been previously well coated inside with glycerin and stoppered at one end with a close-fitting cork. This piece of tubing was then placed in the centrifuge while hot and the centrifuge started rather rapidly. After about a minute of rapid spinning the centrifuge was slowed down and kept going at a moderate rate until the paraffin was completely cooled. Experience has shown that if the small objects are simply allowed to settle and the tube then cooled in water, the paraffin will solidify first next to the glass and remain fluid in the center with the result that the core of the plug will be hollow when the final contraction and cooling has taken place. The hollow may extend quite down to the cork, making it impossible to cut a clean, well-shaped block for sectioning. The use of the centrifuge completely overcomes this trouble and also crowds the minute objects closer to the bottom of the tube against the cork. When the paraffin has cooled, the cork is pulled out and the paraffin plug pushed out of the tube.

The paraffin method was used exclusively for embedding bark and leaf tissues. It was found that the bark tissue could be easily cut down to

with 50 per cent hydrofluoric acid and then embedded in celloidin,¹ or was simply placed in 95 per cent alcohol and glycerin, equal parts, for about 10 days to 2 weeks, and sectioned without embedding at all. Neither of these two methods gave better sections than could be obtained from fresh tissue with the aid of the ether freezing microtome and a sharp knife, except in the case of tangential sections. A thickness of 10 to 20 μ was found the most favorable for the study of the mycelium in the different elements of the cortex, phloem, and xylem. In sections thinner than 10 μ the mycelial strands were often torn out of place. For the study of the pycnia and æcia sections were used as thin as 3 μ , but 5 to 7.5 μ were generally employed. Sections of uredinia and telia were cut 3, 5, and 7 μ thick.

For mounting, Land's fixative,² potassium-bichromate + gum-arabic, was found superior to egg albumen and was used almost entirely for bark sections. It was also found more satisfactory for long sections of the leaf.

A rather long series of stains was employed, and a comparison of the different features of the mycelium and host cells made from slides colored with the different stains. The diagnostic method has already been mentioned. Alcoholic safranin and clove-oil + gentian-violet were particularly good for mycelium in the xylem. Safranin and Delafield's hematoxylin was also a favorable combination for such infected tissue. In the phloem and cortex the mycelium was well differentiated with Delafield's hematoxylin followed by erythrosin in 70 per cent alcohol. For cytological study Haidenhain's iron-alum hematoxylin in combination with aqueous Congo red, aqueous orange G, or *Lichtgrün* in 95 per cent alcohol proved excellent. Host and parasite cell walls stained well with the *Lichtgrün*, whether in combination with the hematoxylin or safranin. Flemming's triple stain³ gave the best results for cytological study, with one or two exceptions to be mentioned later; and by using a strong violet stain the mycelial and host relations at the bases of the sori were brought out more clearly than with any other combination. The gentian-violet was made up in small quantities, enough for one week, as it does not keep well. All reagents from the safranin to the xylol were handled with pipettes from dropper bottles and used but once.

It was found that material once correctly embedded in paraffin, sectioned, and mounted, would stand the jump from 95 per cent alcohol to water, then a 5-minute immersion in full strength hydrogen peroxid for bleaching, without shrinkage or distortion. The sections were rinsed in

¹ See FLOWMAN, ARTHUR B. THE CELLOIDIN METHOD WITH HARD TISSUES. *IN BOT. GAZ.*, V. 37, NO. 6, P. 495-501. 1904.

² See CHAMBERLAIN, C. J. METHODS IN PLANT HISTOLOGY. 63, 2, 314 p. illus. Chicago, 1915.

³ The stains were made up according to the following directions. SAFRANIN. Solution I: 1 gm. of safranin in 95 cc. of 95 per cent alcohol; filter. Solution II: 1 cc. of aniline oil in 95 cc. of distilled water. Shake the two solutions together and filter. GENTIAN-VIOLET. Solution I: 1 gm. of gentian-violet in 15 cc. of 95 per cent alcohol. Solution II: 3 cc. of aniline oil in 80 cc. of distilled water. Mix the two solutions, shake, and filter. ORANGE G: 1 gm. of orange G. in 100 cc. of distilled water.

water after bleaching and kept in 70 per cent alcohol until they could be stained. They were stained in safranin for one to two minutes, rinsed in water, differentiated to the proper point with 50 per cent acid alcohol and washed with fresh 50 per cent alcohol. They were then stained for 15 to 60 seconds in gentian-violet. The violet was followed at once by the orange G with the slide held at an angle so that the orange G would run off rapidly and carry the violet with it. Then a few drops of absolute alcohol were mixed with the orange G which remained on the slide to hasten the removal of excess violet, and the mixture was quickly poured off. The mixing process and the pouring off of the mixture were carried out as rapidly as it was possible to manipulate the reagents and the slide. Then a few drops of fresh absolute alcohol were flowed quickly over the slide. The absolute alcohol was replaced by clove oil, the clove oil by xylol, and the mount then finished in balsam. The whole staining and finishing process consumed only a few minutes and the results were far superior to any obtained by following schedules calling for longer staining. The success of the method depends on having good material properly fixed, embedded, and sectioned, and on the use of fresh reagents. The saving in time and the excellence of the finished slides more than make up for the possible excess use of the more expensive alcohols and oils.

The method of using the iron-alum hematoxylin was also shortened by cutting the time in the mordant by from 10 to 30 minutes, and the time in the stain by from 5 to 30 minutes. The results were particularly good for the study of the centrosomes and dividing nuclei. It is obvious that the shorter methods outlined above have distinct advantages over the longer methods generally advocated, provided they yield satisfactory finished products. The short method¹ for the use of Flemming's triple stain is recommended to investigators who have had trouble with this more or less capricious combination.

MORPHOLOGY

INFECTION OF PINE AND EXTERIOR INDICATIONS OF PRESENCE OF PARASITE PREVIOUS TO SPORE FORMATION

The actual process of the entrance of the sporidial germ tube in infecting the pine has not been observed, and indeed the determination of the actual point of natural infection is an exceedingly difficult problem. Practically all infections first become evident by the etiolation or swelling, or both, of the bark at the node or at some point in the internode. The etiolation may be quite marked, as shown in Plate 48, A, representing a recent nodal infection. Internodal infections most often appear to originate at the base of a leaf fascicle. A striking case of this type of

¹ This short method, varied slightly according to the material to be stained, has been very successfully used at the University of Wisconsin and Columbia University, and the author makes no claim for its originality.

infection is shown in Plate 48, B, a. McCubbin (29) reports that out of 1,007 pine infections examined 925 originated at the bases of leaf fascicles, 14 originated in wounds, and 68 were indeterminate. These conclusions controvert the scarcely tenable theory of Marchal (35) that infection generally takes place through wounds. Observations on a large number of infections at Kittery Point, Maine, indicate that most infections can be classed as either nodal or internodal. However, it should be borne in mind that infection probably takes place very often in the bud while the nodal and internodal cells are scarcely differentiated. Therefore the point at which infection appears to originate after the tissues in the bud have elongated may be either nodal or internodal, depending entirely on the chance growth of the sporidial germ tube. In discussing *Peridermium pini* Pers., parasitic on *Pinus sylvestris* L., Hartig (19) states that infections with this rust also originate either at the node or the base of the leaf fascicle.

The swelling and etiolation of the bark noted above are the two most prominent indications in early stages after infection of the presence of the parasite. The swelling is confined to the bark alone, the wood actually becoming constricted in old infections, and is due to the fact that the mycelium of the parasite forces apart the phloem and cortex cells of the host. No evidence of any increased division has been observed in the bark cells, and there seems to be no stimulation toward gall formation, such as occurs in other cases—for example, on *Pinus virginiana* Mill. under the attack of *Peridermium cerebrum* Peck. In the case of the nodal infection in Plate 48, A, it will be noted that there is little or no swelling; but, as stated above, both swelling and etiolation may occur at the same time. The irregular edge of the etiolated area marks fairly definitely the advancing tips of the invading hyphae, which generally extend a little beyond the line.

Other less common external indications of the parasite are bunched needles on seedlings or transplants and occasionally on older trees, and short adventitious branches which spring from the infected nodes. It frequently happens that certain secondary fungus parasites become established on the area already attacked by the rust, and suppress the latter so completely that spore production is partly or completely inhibited. Under these conditions the bark dies and shrinks more rapidly than under the attack of the rust alone, and the stem then appears to be constricted, more or less irregularly, at the infected part. All of these external indications are valuable aids to early recognition of *Cronartium ribicola*.

The period of incubation, from the time of infection to spore production, varies from one to several years, depending possibly on whether the infection takes place in early summer, and therefore under favorable conditions for growth before winter, or in early fall, when growth has unquestionably slowed down. The growth of the mycelium in the bark

in some cases seems to vary directly as the succulence of the host. It is at this time impossible to determine what part weather conditions play, although it seems evident that, once infection has taken place, external conditions cause little or no change in the growth of mycelium. The time of production of æcia in the spring varies directly with warm and favorable weather conditions.

MYCELIUM IN PINUS STROBUS

A brief summary of the main characteristics of the mycelium and the relation of the parasite to the cells of the cortex and phloem of the pine host has already been published (7); however, both subjects will bear review in greater detail. No data can be given at this time on the manner in which the mycelium enters the host tissue, because the youngest stages of infection have not been observed. The course of the hyphæ is always intercellular (Pl. 53). Conditions found at the edge of the infected area indicate that the fungus makes its advance first in the most recent phloem parenchyma and in the rays. This holds true for infections of several years' standing, as well as for those which, to judge from the size of the diseased area, could not have been more than a year old. The hyphæ are relatively large, 3 to 5.5 μ in diameter, and are divided by cross walls into cells which vary considerably in length (Pl. 53). Each cell has a single conspicuous nucleus. In transverse and radial sections the hyphæ can be traced along the rays past the cambium into the xylem (Pl. 53). Here they are confined to the region of the ray cells, with the exceptions noted below. Because of difficulties in cutting transverse sections, which are at right angles not only to the "grain" of the phloem but also to the general course of the hyphæ, such sections are not favorable for study, except in comparison with other sections cut tangentially or radially. In tangential longitudinal sections of the cortex and phloem the mycelium is frequently found in strands, which are especially prominent in the outer phloem region. The strands may surround groups of the phloem parenchyma. In some cases the older sieve cells appear to be filled with hyphæ; but careful observation shows that they have been split apart and flattened out by the hyphæ, which have forced their way into the enlarged intercellular space, and thus practically occupied the same amount of space that was formerly occupied by the living sieve cells. In the rays the hyphæ frequently fill the intercellular spaces adjacent to the horizontal walls of the ray cells, and haustoria from these hyphæ penetrate the ray cells. Resin cells are also penetrated by haustoria. In the xylem the same general conditions are met with in respect to the ray cells that are found in the phloem. Short haustoria arising from the hyphæ in the angles between these ray cells bore through the thin walls of the adjacent tracheids and enter the lumens

of the tracheids¹ for a short distance (Pl. 49, C). Such haustoria are naturally limited in development.

Radial longitudinal sections are more easily cut from bark or wood than transverse or tangential longitudinal sections because radial cuts are splitting cuts. Moreover, in radial sections the vertical widths of the rays are exposed, and therefore all the infected cells of the rays, with the haustoria they contain, as well as the adjacent infected phloem and xylem cells, are more readily examined (Pl. 53). The hyphae lying along one phloem ray frequently are united with similar hyphae lying along adjacent rays by connecting strands. The latter may work their way between phloem parenchyma or sieve tubes. In the xylem the ray hyphae may be connected in the same way by hyphae, which pass from one ray to the other in the intercellular spaces between the tracheids. Often the edge of the ray is bordered by a hypha lying in the space between the outer ends of the marginal cells of the ray and the tracheids (Pl. 53). In the resin-duct parenchyma and the cells lining the duct all the cells are usually penetrated by haustoria (Pl. 53). This applies to vertical ducts in the wood and bark and to horizontal ducts in fusion with rays.

The general features of the mycelium such as their more or less uniform diameter, with occasional bulges where there is a little extra room in an intercellular space, their relatively large size, uninucleate cells, and constant relations to the host cells of the different tissues, are characteristic enough to make the mycelium alone a sufficient basis for the recognition of *Cronartium ribicola* before any spores are produced. The haustoria often the most striking objects in the infected cells, are the most important elements of the mycelium from a diagnostic standpoint. The haustoria apparently have the power to pierce the cell wall at any point (Pl. 53). Young haustoria are usually straight (Pl. 58, B), constricted at the point of passage through the wall, and irregularly bulging inside the cell (Pl. 53; 58, C, D). Their outline soon becomes more or less irregular. In the phloem parenchyma they do not reach the development found in the ray cells, where they coil on a wide spiral or curve at rather sharp angles. The curve and the spiral are probably different expressions of the same process—that is, the adjustment between the growing haustorium and the cytoplasm of the host cell. As Sappin-Trouffy (50, 51) has pointed out, the haustoria of the rusts appear to seek out the nucleus of the host cell, and sometimes even entwine it. No such extended development has been observed in the case of the haustoria under discussion, but it frequently happens that the host nucleus is deated (Pl. 53) by the tip of the haustorium. Olive (42) shows that the haustoria of *Botryorhiza hippocrateae* Whetzel and Olive form botryose masses which may almost completely fill the host cell. In this fungus the haustoria

¹ In a previous article (7) the statement was made that haustoria do not enter the wood cells. Improved technique has shown this statement to be an error.

are multinucleate; in *C. ribicola* they are apparently always uninucleate (Pl. 53). In the latter fungus no case of more than one cell in a haustorium has been observed by the writer, although as many as three tips are not uncommonly seen. It is difficult to determine whether the haustorium is always cut off from the hypha from which it is originated and this point will have to be left undecided.

When properly stained with any of the combinations given on page 623, a sheath can be made out at the base of each haustorium (Pl. 58, C, D), enveloping it for a distance of several microns, as if the haustorium were set in a cuplike holder. The sheath really extends all over the haustorium, but is generally very thin in the middle region. At the tip it is as thick or thicker than at the base (Pl. 58, E). In its staining reaction the sheath resembles the wall of the host cell. There is at least a possibility that it is formed by the host cytoplasm in response to the irritation or stimulus caused by the presence of the haustorium. Olive (42) states that the plasma membrane of the host protoplasm is pushed in by the haustorium of *Botryorhiza hippocrateae* as it invades the host cell. Apparently the cytoplasm shrinks away from the advancing haustorium in some cases, leaving an appreciable space. This phenomenon has not been observed in *Cronartium ribicola*, but it is certain that the plasma membrane of the host cell is not broken nor pierced by the haustorium; it must be pushed in as the tip of the haustorium grows. The greatest points of irritation produced by the haustorium in the host cell would be the point in the cell wall through which the haustorium entered and the point of contact of the advancing haustorial tip with the host cytoplasm. Consequently it might be assumed that the greatest results from the irritation would be observed at these two points—namely, at the base of the haustorium and at its tip. At these two points the sheath is thicker than at others. Possibly the narrow space which must occur between the haustorium and the host cytoplasm membrane is the dumping ground for precipitation products resulting from the irritation. These products might constitute the beginning of the sheath which would gradually increase in thickness as the age of the haustorium increased.

The sheath certainly does not come into being at the time the haustorium penetrates the wall, as Smith (53) has reported for the haustorial sheaths of the Erysipheae. It has not been found in connection with young haustoria in any case and seems to be an accompaniment of maturity or old age. The hole in the host cell wall, through which the young haustorium passes, is comparatively small (Pl. 58, C, D) and at first the wall is no thicker at this point than at any other. Whatever thickening takes place at the point of penetration occurs after the haustorium has entered the cell.

The young haustorium is full of cytoplasm, with a typical round nucleus (Pl. 58, B). As the wall of the haustorium, which is at first

quite thin, thickens and the sheath develops, the cytoplasm becomes vacuolate (Pl. 58, C, D, E), and the nucleus loses its normal structure, condensing into a shrunken deeply staining mass. In such a state the haustorium can not function efficiently as an absorbent organ. At any rate the host cell and its nucleus do not show much evidence that the parasite seriously interferes with the normal cell growth. Even when the cells are completely separated by mycelial stands, the host nuclei may remain apparently normal in shape and structure.

PYCNIA

On any given area of infection the pycnia precede the aecia, by at least one growing season. Succeeding generations of pycnia and aecia follow a more or less definite schedule. The plan of the advance of the fungus is illustrated in Plate 54, A. Immediately surrounding the point of infection the bark may show pycnia and aecia at the same time (Pl. 48, B), but in such cases the pycnia have either passed maturity or the aecia have developed abnormally early. After the infection has spread for some distance, provided that the tree is not less than several inches in diameter, the pycnial area is normally always in advance of the aecial area. The sequence of the etiolated bark, pycnial area, and aecial area is evident in Plate 54, A.

Very young pycnia are difficult to find. They seem to develop rapidly when once started, for none have been examined which were mature in the center and still young at the periphery, a condition which is commonly met in aecia and uredinia. The hyphae which contribute to formation of the pycnium force their way between the cells of the outer cortex in a direction at right angles to the outer surface of the bark (Pl. 49, B, a). They are aggregated between the outermost cortex cells and the periderm, forming a layer of pseudoparenchyma two to four cells thick (Pl. 49, B, b). From this layer arises a series of short branching trunks (Pl. 58, A, c). Each of the ultimate branches from these trunks is a long sporophore, on the tip of which a number of pycniospores are formed, one after another (Pl. 49, A, B; 58, A, a, b). Roughly speaking, the pseudoparenchyma makes up one-fourth, the short branching trunks one-fourth, and the sporophores one-half of the vertical width of the sorus (Pl. 49, B, 58, A). Pycniospores are produced in large numbers. Mixed with a thin sweet gelatinous fluid, they collect between the sporophore tips and the periderm layer (Pl. 49, A) forcing the latter up into the form of a shallow blister. Finally a small break in the periderm layer allows the spores to escape, together with the sweet fluid, in the form of a honey-colored drop called the pycnial drop (Pl. 48). The pycniospores are typically pyriform (Pl. 58, A, b), measuring when mature approximately 2.5 by 3.5 μ . Their mode of formation appears to agree closely with that of the pycniospores in other rusts. At the rounded tip of the sporophore a bud is formed which swells until it reaches the size of a mature spore

(Pl. 58, A). After receiving its nucleus the spore is abstricted. No evidence of a collar like that figured by Blackman (2) was seen, unless the odd constriction of the cytoplasm shown in Plate 58, A, c, can be considered as a collar. This phenomenon is quite common. Occasionally long hyphal filaments grow some distance out beyond the tips of the sporophores. The structure of the spores is typical of pycniospores in general. They appear to be completely nonfunctional. No attempt was made to germinate them.

The dark areas on the bark which indicate the location of the pycnia are designated pycnial spots (Pl. 54, A, b; 48, B). They are honey-yellow to brown-yellow at first, but they gradually assume a color like that of clotted blood as the pycniospores mature and ooze out, and may finally become almost black. Later, when the covering of cork cells sloughs off or is eaten off by insects, the drying cells of the host tissue beneath turn a typical light pink color. After reaching maturity the pycnium is cut out from the host tissue by the formation of a cork cambium and the deposition of a cork layer at a depth of four to six cells below the bottom of the pseudoparenchyma base of the sorus (Pl. 50, C, b). It is the exposure of this layer which reveals the typical color of newly formed cork cells. The pycnial spots, whether young or old, are valuable diagnostic characters, because they often make possible the detection of infected trees before æcia are produced—that is, in time to destroy such trees before æciospores can spread the disease.

ÆCIA

Æcia appear in April, May, and June. Very often the whole area on which æcia can normally develop in a given season is covered with closely crowded sori pushing their way through widening cracks in the bark. By the middle of May the peridia are usually broken and the spores escape in orange-yellow pollen-like showers. Spore production continues for some time after the æcia open. Young æcia are easily obtainable at the edge of the æcial area where their presence is indicated by a thin yellow line just beneath the outermost layers of the bark cells. The hyphæ contributing to the formation of the æcium are aggregated into a mycelium, which is clearly made up of elements running tangentially among the host cells at a depth of 6 to 10 cells below the periderm layer. By their continued growth these hyphæ force the host cells apart, so that the latter become isolated and embedded in a matrix of tangled mycelium (Pl. 50, B). This separation may extend to a depth of 15 to 20 cells in the cortex. About 6 to 8 cells below the periderm layer a mass of pseudoparenchyma is formed by the packing together of the hyphæ. In the pseudoparenchyma a layer of fertile cells becomes discernible by their denser protoplasmic content (Pl. 50, A, b; 54, B, f*c*).

The fertile cells cut off rows of sterile cells (Pl. 54, B, s*c*), 6 to 12 cells long, which may increase in size at first, but which later degenerate to make

room for the developing aëciospore chains. These sterile cells correspond to the buffer cells of Olive (41) and Fromme (14, 15). The cells of the fertile layer, which are somewhat larger than those of the vegetative mycelium, may fuse in pairs, the fusion beginning in the center of the sorus and proceeding centrifugally toward the periphery, as is normally the case in the aëcium and its analogs. As a result of the fusion and union of the cell contents of the contributing cells a large cell is formed, the basal cell (Pl. 58, I, J), which will give rise acropetally, by repeated division, to the aëciospore chain. The fusion seems to be complete as far as the cells are concerned, and evidently takes place rapidly, as there is only occasional evidence even in very young aëcia of remaining parts of the cell walls which originally separated them (Pl. 58, I). The two cells are not always at the same stage of development, as one of the cells is often shorter or at a lower level than the other (Pl. 58, J). However, there is little ground for considering the two cells as different in kind. The essential process is the cell fusion and consequent association of the nuclei to form the dikaryon¹.

Fusion of more than two cells also occurs, trinucleate and tetranucleate basal cells being quite common. Aëciospore chains arising from such basal cells may contain the same number of nuclei as the basal cells, just as in aëcia of other genera. Trinucleate basal cells are very numerous in young sori and at the edge of older ones, but the number of trinucleate aëciospore chains is considerably less than one would be led to expect from the number of the trinucleate basal cells. There is a possibility that the extra nucleus frequently degenerates, evidence of this being occasionally seen. This point will be discussed briefly later on. Multiple cell fusions of a more complex character are also common (Pl. 58, K, L) recalling Olive's (41) and Fromme's (14) observations. The nuclei and cytoplasm of a number of adjacent cells, not only those in the upper row of the fertile layer but also others considerably below that level, become associated in one large irregular cell (Pl. 58, L). What happens to these large cells is not clear. The probability that the multiple fusion cells may give rise to a number of spore chains is reservedly suggested here.

FORMATION OF PERIDIAL CELLS AND AËCIOSPORES

The basal cell divides into an upper part, the aëciospore initial cell, and a lower part, potentially equivalent to the primary basal cell (Pl. 58, M). Each of the first few cells cut off from the basal cells normally divides into two cells of unequal size. The larger cells thus formed adhere more or less completely into a layer three to five cells thick which constitutes

¹ The term "dikaryon" is to be preferred to that of "syngkaryon," the more common term, because of the earlier use of the latter name to designate the fusion nucleus resulting from the union of the male and female gamete nuclei by zoologists. Pavillard's (43) objection to syngkaryon should be sustained, and Maire's (32, 34) suggestion that dikaryon be substituted for the older word has the virtue of correcting an error and supplying a distinctive term for the unique condition found in the rust sporophyte and the Basidiomycetes in general.

the peridium (Pl. 56, B); the smaller ones go to pieces. At the periphery of the sorus several adjacent basal cells cut off units which never become functional aëciospores but which always form peridium. The small cells between these peridial cells indicate that the potential aëciospore initial, although destined to become a peridial cell, regularly divides to form two cells homologous to an aëciospore and an intercalary cell (Pl. 56, A). Thus, the multiple-layered peridium of *Cronartium ribicola* is formed in the same general manner as reported by Fromme (15) for the peridia of other deep-seated aëcia. When first formed, the individual cells are subspherical to elliptical, and smooth-walled. As the sorus matures, their walls thicken and their outline changes according to their position in the peridium. All tend to become more or less polyhedral. The cells at the top of the sorus are usually more rounded than those at the sides, since the latter are elongated by the pull exerted on the sides as the developing spore chains force the central part of the peridium out. Therefore, the size of the cells varies widely (18 to 40 by 12 to 42 μ). The mature wall is 3 to 9 μ thick. The walls of the outermost cells are smooth or slightly granular, while the inner cell walls are studded with short tubercles which sometimes appear to mesh with those of the adjacent cells, perhaps contributing thereby to the strength of the peridium as a whole (Pl. 56, B). The cell contents of the peridial cells slowly degenerates until they become empty shells.

After the cells which form the peridium are abstricted from the basal cells, the aëciospore initials are cut off. Each aëciospore initial cell undergoes division into a larger upper cell, the aëciospore, and a smaller lower cell, the intercalary cell (Pl. 58, W, M, b, X, Y, c, d). By the repeated divisions of the basal cell a row of alternating spores and intercalary cells is formed which constitutes the aëciospore chain (Pl. 50, B; Pl. 58, Y). After the division of the aëciospore initial cell, both resulting cells, the aëciospore and intercalary cell, grow rapidly. The aëciospore reaches its normal broadly elliptical shape when about three or four spores distant from the basal cell. The intercalary cells elongate, eventually becoming mere thin connecting elements between the aëciospores in the chain, and finally disintegrate entirely. The spore wall thickens greatly when the spore has attained its full size. A thin space in the wall, suggesting the germ pore of other rust spores, is evident at the point of attachment of the aëciospore and the intercalary cell below it (Pl. 58, Y, Z). This thinner place in the wall may persist even in the completely matured spore. It is comparable to similar phenomena in other rusts and does not normally function as a germ pore. The mature spore measures 18 to 21 by 20 to 26 μ . The aëciospore wall seems to be made up of two parts, an endospore overlain by a somewhat thicker exospore. The latter is distinctly characteristic of aëciospores of *Peridermium* spp. Part of it is cracked up into tubercles or warts, which makes it decid-

area is more or less indefinite in extent at the basal end of the spore. Verrucose and smooth areas grade one into the other along the irregular line which separates them. The smooth area is fissured near its edge into blocks which become smaller and smaller until they approach the size of the tubercles of the verrucose area (Pl. 58, Z).

No completely satisfactory explanation of the manner in which this type of spore sculpturing arises has come to the writer's attention. The following theory is reservedly offered. The two walls of the spore are present when the spore is quite young, and both continue to grow and thicken up to a certain point or until the spore has nearly reached its full size. The outer wall hardens more rapidly than the inner one and in consequence becomes fissured irregularly as the still elastic inner wall continues to expand under the pressure of the growing spore content. The longer growth in size continues the more complete will be the fracturing process and the larger the verrucose area in proportion to the smooth area. Experimentally, the smooth area can be converted into verrucose area by soaking the spores in water. After several hours, if germination does not take place in the meantime, the spore absorbs water enough to cause it to increase appreciably in size. The tubercles of the verrucose area become free from their attachment to the inner wall and float around in the water. The smooth area, under the expansion pressure exerted as the inner wall swells, cracks and fissures until it becomes irregularly verrucose, approaching the conditions found on the normal verrucose area of the mature spore. The process of fissuring can be watched quite easily. The experiment at least suggests the manner in which a type of spore sculpture so oddly irregular could arise.

The spore wall on the smooth area has been heretofore considered to be thicker in section than the verrucose area. Examination of complete sections of the spores shows that this is not always the case. In fact, it is only in occasional instances that it holds true. As a rule, there is no appreciable difference in the thickness from the inner edge of the wall to the outer tip of any given tubercle and the thickness from the inner edge of the wall to the outer edge of the smooth area. When dealing with whole spores, refraction phenomena increase the difficulties in measuring the true thickness of a curving wall of the type presented in the smooth area of the æciospore of *Cronartium ribicola* and may account in part for the misinterpretation of the actual condition.

GERMINATION OF ÆCIOSPORES

Germination of the æciospores may take place rapidly under favorable conditions, but as Maire (32) aptly puts it "*la germination des æcidiospores * * * est parfois très capricieuse.*" The work of Spaulding and his assistants has shown that spores which would not germinate at all in hanging-drop cultures, on a water film, or on moist filter paper, either at room temperature or in the ice box, or at room temperature after cooling

in the ice box, were perfectly capable of producing infection on species of *Ribes*. These results confirm those of Klebahn (25), who found that aëciospores which would not germinate in water did germinate very rapidly on leaves of *Ribes* spp. and only less rapidly but still abundantly on a gelatinous decoction of leaves of *Ribes* spp. These experiments suggest the probability that some direct chemotactic stimulus is exerted by the leaves of *Ribes* spp. on the aëciospores. Klebahn has pointed out that there is considerable difference between the faculty for germination, as determined by artificial cultures, and the faculty for infection. Spatlding and his assistants have further determined that aëciospores frequently germinate—even then only a relatively low percentage of fresh spores do so—more readily after cooling in the ice box than at room temperature, and that sometimes they have to remain in the ice box to secure germination. Too much water is often as inimical to germination as too little. A single spore may produce one to several germ tubes (Pl. 59, A), which attain considerable growth in artificial cultures. Where the germ tube passes through the heavy exospore it is constricted as shown in Plate 59, B. The tubes branch freely. The protoplasm is densest at the advancing tips of the hyphae.

INFECTION OF RIBES SPP. AND MYCELIUM IN THE LEAF

Whether the germ tubes have the power to pierce the upper epidermis of the leaf of *Ribes* spp. or must always come to rest in a favorable position on the lower epidermis in order to cause infection is not definitely known. All the evidence gathered from the examination of artificially inoculated leaves points to the conclusion that infection occurs normally as a result of the germination of the aëciospore on the lower surface of the leaf and the subsequent passage of the germ tube through a stoma. No evidence either of any break in the upper or lower epidermal cells, or of the remnants of any hypha passing through them, has been discovered. Furthermore, the mycelium is always abundant in the air chambers adjacent to the stomata, even in very young infections, and occasional remnants of spores and hyphae near and in the stomata point to the stomata as the avenue of infection.

The first indication of infection in the leaf of species of *Ribes* is often indicated by the paling of the infected areas. Sections of such areas show that the mycelium has spread in the intercellular spaces and air chambers of the mesophyll. Haustoria (Pl. 59, D) enter all types of the leaf cells, with the possible exception of the xylem elements of the bundles, although they are comparatively rare in the epidermal cells. The cells of the mycelium and the haustoria are binucleate (Pl. 59, C, D). Generally there is a much smaller relative amount of mycelium in the leaf tissue than among the same number of host cells of the pine. The loose structure of the mesophyll allows plenty of room for the hyphae to grow without severe crowding of the host cells. In fact, the hyphae are aggregated only at the time of production of uredinia or telia.

UREDINIA

The development of the uredinium is illustrated in Plate 51, A, B, and 55, A-C, which form a series from the very beginning of the formation of the sorus to its maturity. When the uredinium starts to form, the fungus cells may be found aggregated in groups in some large air space, generally near a stoma. Certain of the cells in each group become oriented with their long axis more or less at right angles to the epidermis, against which they are closely appressed (Pl. 55, A). These cells are functionally equivalent to the basal cells of the mature sorus, in that from them arise the cells which go to make up the peridium, and the first urediniospores. The first division of any one of the vertically elongated cells results in the formation of a cell which adheres to its neighboring homologous cells to make the peridium (Pl. 55, B, a). In the second division the first urediniospore initial cell, or its equivalent, is cut off. This divides immediately to form the first urediniospore and stalk cell or their equivalents. The position of the cells referred to, as they appear in a young uredinium, are illustrated in Plate 55, B, in which *a* is the peridial cell, *b* the young urediniospore or its homolog, *c* the stalk cell or its homolog, and *d* the basal cell. It will be noted that the arrangement of the cells suggests that the sorus is made up of a compact aggregation of vertical rows of cells. This arrangement is temporary for the middle region of the sorus, but permanent for the circumference of the sorus. Plate 55, B, which represents a median section from a young uredinium, can be duplicated by taking sections through the edge of any mature sorus. This fact should be kept clearly in mind in any discussion of the structure of the uredinium.

The first urediniospores mature in the middle of the sorus (Pl. 51, B, b). Their formation and method of growth corresponds closely with the production of normal stalked urediniospores in other rusts. To repeat the process suggested above: The basal cell undergoes division (Pl. 55, C, *c*; 59, E-I). The upper cell is the urediniospore initial (Pl. 59, J, L); the lower cell is potentially the equivalent of the original basal cell. The urediniospore initial now divides (Pl. 59, K, L, M) to form a larger upper cell, the urediniospore, and a smaller lower cell, the stalk cell (Pl. 51, B, *b*, *c*; 55, C, *d*). While this process is going on, the layer of cells constituting the peridium gradually separates from the underlying urediniospores along the line between the cells marked "*a*" and "*b*" in Plate 55, A. The figures in Plate 51, A and B, represent steps in the process leading up to conditions shown in Plate 55, C. As the urediniospores grow, the peridium (Pl. 51, A, *a*; B, *a*) is forced up into a dome. The individual peridial cells lose the regular shape and outline shown in Plate 59, B, *a*, and become irregularly compressed or obliquely flattened. The growing urediniospores develop pressure against the epidermal cells, which flattens them out and finally causes them to be torn apart (Cf. Pl. 51, A, B, with Pl. 55, C). The break in the epidermis frequently

comes at a stoma, sometimes extending from one stoma to another; but the break in the peridium, when it finally occurs, is confined more or less definitely to the top of the dome. Usually the peridial cells around the break are irregularly thickened (Pl. 55, C). The orange-yellow urediniospores work their way out through the opening, sticking more or less closely to each other to form a spore crown on the top of the peridium. The basal cells continue to cut off urediniospore initials by a process similar to that involved in the formation of the primary initials, which has been described above. The secondary urediniospore initial may be formed alongside the base of the stalk of the first-formed urediniospore before the latter has reached maturity (Pl. 55, C, c; 59, N). The secondary initial then divides to form a urediniospore and stalk cell, in the same way as the primary initials. Meanwhile the stalk cell of the primary urediniospore elongates, withers, and goes to pieces. While no more than two spores have been found in connection with a single basal cell, each basal cell must frequently give rise to several spores, to judge from the number produced in a single sorus. The size of the mature spore (Pl. 59, P) is 10 to 20 by 19 to 45 μ .

The spore-bearing basal cells are confined to the middle part of the sorus. In figure C of Plate 55 it will be noted that on either side of the group of spores there is a group of parenchyma-like cells (*g*) made up of units which are arranged in more or less vertical rows. The cells of any individual row of four cells may be homologized with the cells lettered *a*, *b*, *c*, and *d* in figure B of Plate 55. This parenchyma-like tissue forms an encircling bank of cells which completely surrounds the mature uredinium. In this tissue the row arrangement of the cells shown in figure B persists, although the cells themselves lose their contents and become practically dead by the time the basal cells in the middle of the sorus are actively forming urediniospores.

The uredinium seems to be limited more definitely than either the pycnium or æcium in its ability to extend in a centrifugal direction. Its extent is predetermined, much more exactly than in the case of the two other sori, by the amount of massed mycelium from which it arises. Both pycnia and æcia originate in tissue which is comparatively full of mycelium before they start to develop. In the leaf of species of *Ribes* there is rarely any massing of the hyphæ to form the packed mycelium so common in the pine host, and the leaf cells are only occasionally distorted except in the immediate vicinity of the uredinia and telia.

GERMINATION OF THE UREDINIOSPORES

Urediniospores exhibit the same irregularity in germinating in artificial cultures as do the æciospores. Even when they are dusted over the surface of young, fresh, moistened leaves and placed in a damp chamber they may or may not germinate in large numbers. On a water film they absorb water and swell considerably. The cytoplasm becomes vacuo-

late. The germ tube passes through the exospore without the aid of a germ pore, assuming at first the shape of a more or less swollen vesicle (Pl. 59, Q). This lengthens rapidly into the young germ tube (Pl. 59, R), into which pass the contents of the spore. The endochrome and protoplasm are densest at the tips of the tube and in its branches. Duff (12) has shown that change of temperature stimulates the urediniospores to germinate and that light, especially ultra-violet light, may completely inhibit germination. This action of light may have a direct bearing on the apparent failure of the spores to cause infection at long distances from their source. All the evidence at hand points to the conclusion that the germ tube causes infection by passing through the stomata on the lower surface of the leaves. However, the tube may extend some distance over the leaf surface before actually entering a stoma, in such cases passing directly over stomata in its path. Urediniospores may retain their viability for some weeks. The uredinium is the repeating sorus in the life cycle of *Cronartium ribicola* (fig. 1), and the spores in succeeding generations infect leaves of *Ribes* spp. until late summer.

TELIA

Telial columns arise either from old uredinia or as entirely new and separate entities. They appear later than the uredinia and are more common in late summer. In the fall they are usually the predominant spore generation present. In the greenhouse they are produced throughout the year. On a given infected area the columns may occupy the central part where the uredinia were first produced, surrounded by a narrow peripheral region bearing the most recently formed uredinia. The development of the telial column is the same whether it is from an old uredinium or in a new sorus and, as the latter is probably the most common occurrence, it will be described. The massing of the hyphae, formation of the peridium, and the parenchyma-like cells which surround the spore-bearing part of the sorus, and the orientation of the basal cells proceed exactly as in the uredinium. It is impossible to tell very young uredinia and telia apart. The binucleate basal cells undergo division (Pl. 59, S, T, U, V, W) in the same manner as in the case of the uredinium, but the cells cut off do not divide as in the other sori described. Instead they lengthen out and become teliospores. Each basal cell cuts off a vertical row of spores (Pl. 52, A, B), the central cells of the sorus producing spores slightly ahead of the cells at the periphery, as in the æcium and uredinium. As the spore columns lengthen, the peridium is pushed up into a dome (Pl. 52, A), which later ruptures (Pl. 52, B) and goes to pieces. The spores lengthen and soon reach full size, at about which time they become provided with a substantial wall slightly thickened at the upper end (Pl. 57, A, B, C). The first spores cut off—those at the tip of the column—are more or less polyhedral (Pl. 52, B; 57, A); the other spores are typically broad spindle-shaped (Pl. 52, D;

57, B). They usually touch, but do not crowd one another. Adjacent to the points where one spore abuts on its neighbors there may be considerable space. Cross sections of the columns (Pl. 52, E) show that the spores are only very slightly angular as a result of mutual pressure. Although they vary widely, normal mature teliospores average approximately 16 by 42 μ , with a wall about 2 μ thick. They are uninucleate, as a result of the fusion of the two nuclei which are normally present in the young teliospores. The mature telial column is an aggregation of a number of vertical rows of mature teliospores (Pl. 52, C). The length and diameter of the columns vary greatly; they may attain a length of 2 mm. and an average width of approximately 100 μ . They are usually curved or semispiraled, a result apparently of unequal development of the spores of some of the rows. They vary from almost spherical to irregular ovoid or elliptical in cross sections. Occasionally abortive, nondeveloped spores are scattered throughout the length of the column (Pl. 52, C, D). Other abnormalities will be discussed briefly later.

GERMINATION OF TELIOSPORES AND PRODUCTION OF SPORIDIA

All of the spores in the telial column may germinate *in situ* (Pl. 56, C). The exospore pushes out at some point in the form of a rounded papilla, which ruptures and allows the extrusion of a stout germ tube—the young basidium or promycelium (Pl. 57, D). This reaches its full size in a few hours and then becomes divided into five cells (Pl. 57, R). From each of the four upper cells arises a stout sterigma, on the tip of which the sporidium swells to its full size (Pl. 57, S–V). When abstricted, the sporidium is almost exactly spherical, measuring approximately 8 to 10 μ in diameter. At one point on the thin wall is a tiny papilla-like swelling, which marks the point of attachment to the sterigma (Pl. 57, AA, EE, GG).

The germ tubes from the spores in the middle of the column work their way out through the intersporal spaces and then develop in the manner described above. If the teliospores germinate under water, the germ tube lengthens out into a narrowly spiraled or twisted hypha; in other words, promycelia do not develop unless they have access to the air. The color of the germinating column, a very pale pink, gives a distinctly characteristic appearance to the fungus at this stage.

GERMINATION OF SPORIDIA

The sporidia germinate in artificial culture by sprouting relatively stout germ tubes (Pl. 57, X, Y, Z, BB) which probably continue growth in the normal fashion under favorable environment. In many cases, however, the short germ tube swells at its tip on reaching a length of several microns and the swelling becomes a secondary sporidium, apparently exactly similar to the primary one (Pl. 57, CC, DD). Sappin-Trouffy (51) has figured the same phenomenon for the sporidia of *Cronartium flaccidum*.

Alb. and Schwein. on *Peonia officinalis*. It is apparently a common occurrence. Just how the sporidia reach the pine is not known, but it may be assumed that they are usually air borne. Infection of the pine host follows their germination under favorable circumstances on young pine shoots.

CYTOLOGY

* PREVIOUS INVESTIGATION

Our knowledge of the cytology of the genus *Cronartium* is very fragmentary; in fact, no reference has been found to the nuclear phenomena accompanying the formation of any bark-inhabiting æcium of this group. Poirault and Raciborski (44) figure silhouettes of the nuclear division, as they interpreted it in the formation of the æciospores of the form known to them as *Peridermium pini acicolum*.¹ Sappin-Trouffy (51) gives a fair diagram of this acicolous type of æcium and the nuclear division at its base. His diagram of the telial column of *Cronartium flaccidum* indicates the phenomena accompanying the production of teliospores, though the figures are too minute to be more than suggestions of the actual conditions. He regards the processes as closely similar to the nuclear phenomena in other rusts, in which he is quite correct. Aside from the work of these authors, nothing beyond incidental mention of the cytology of the genus *Cronartium* has come to the writer's attention.

It is hardly necessary to give an extended resumé of the work of previous investigators on the cytology of the rusts, on account of the excellent reviews which have been published by Blackman (2), Christman (4, 5, 6), Fromme (14, 15), Maire (32), Guillermond (16), and Ramsbottom (46). Moreau (36, 37, 39) revives the older view that there are but two chromosomes in the rust nucleus, which hardly seems tenable, and establishes the presence of centrosomes in the resting nuclei (36). This subject will receive further consideration later. Fromme's (15) recent paper on the morphology and cytology of the æcium clears up the matter of the formation of the deeper seated sori and establishes the similarity of spore formation in the caoma and æcium. The life history of a complete rust is divided into two stages: the gametophyte, with uninucleate cells, and the sporophyte, with binucleate cells. The gametophyte begins with the reduction division, or its equivalent, in the promycelium and continues up to the fusion of the gamete cells at the base of the æcium, which initiates the dikaryon. The sporophyte begins with the inception of the dikaryon and continues up to the reduction division, or its equivalent, in the promycelium. The association of the nuclei in the basal cells of the æcium is regarded as the equivalent of a true fertilization, and the fusion of the two nuclei in the young teliospore as the completion of the process necessary for a mixing of the chromatin elements previous to reduction (31). The nuclear divisions are true mitotic divisions, accom-

¹ Vuillemin (52) also reports some of the phenomena accompanying æciospore formation in *Peridermium pini*.

panied by centrosomes or their equivalent, true spindles, and chromosomes. The character of the latter is still unsettled. In the dikaryon both nuclei divide simultaneously in conjugate division.

The chief differences of opinion have been in regard to the nature of the two cells which fuse to form the basal cells and in regard to the sterile cells. Blackman (2) held that the two basal cells were unequal, either in size or time of development, and that one, the larger, received the nucleus of the other, the smaller, by a process of nuclear migration through a comparatively small pore. After Christman's paper (4) announcing the fusion of two equal cells by the complete absorption of their appressed walls, Blackman and Fraser (3) investigated a number of forms and stated that the dikaryon might arise (a) through the process of nuclear migration from one cell to another, as Blackman first reported, or (b) by a similar nuclear migration from one vegetative cell to another below the fertile layer, or (c) by the process described by Christman. Most of the evidence since brought to view has supported Christman's theory. Olive (41) has pointed out that the manner of fusion is not of great importance, inasmuch as the conjugating cells are not definitely organized as male and female organs; for the essential result of the fusion is the establishment of a dikaryon by the association of two nuclei and their accompanying cytoplasm, regardless of their individual origin.

The sterile cells cut off from the fertile cells before fusion have been regarded as homologous to the trichogyne of the Florideae (2); as buffer cells which protect the developing basal cells below (15, 41); or as degenerate female gametes, which once were fertilized by the now nonfunctional pycniospores. According to this last theory, advanced by Moreau (40), the cells as a group constitute a *preecium* (*preecide*) and the individuals *preeciospores* (*preecidiospores*). Fromme (15) has shown that the number of sterile cells varies considerably in different species and suggests that they are formed in response to the general conditions of environment under which the particular sorus happens to be developed.

In the following presentation of the cytological details in *Cronartium ribicola* the fusion nucleus in the mature teliospore will be taken as the starting point, inasmuch as it is the first stage in the nuclear history of the gametophyte generation. The nuclear processes accompanying the production of the different spore forms can then be described in logical cytological sequence and at the same time in chronological order.

NUCLEAR PHENOMENA IN FORMATION OF PROMYCELIUM AND PRODUCTION OF SPORIDIA

With the germination of the teliospores the morphological history of the gametophyte may be said to begin. Coincident with this germination, changes occur in the fusion nucleus. As the promycelium reaches

its full size the fusion nucleus, formed by the union of the two nuclei in the young teliospore, migrates from the teliospore into the promycelium and starts to divide (Pl. 57, E). The membrane becomes irregular and disappears (Pl. 57, F) at the same time that the chromatin granules increase in size. These granules become longer and more deeply-staining units, which form a tangled mass in the position occupied by the fusion nucleus when it moved out into the promycelium (Pl. 57, G). A definite spindle, with centrosomes at the poles, then develops. At the middle of the spindle the chromatin tangle becomes arranged in an equivalent of the equatorial plate stage. In the next stages the chromatin is in the form of separate units (Pl. 57, H, I), which can be clearly seen under favorable conditions and proper differentiation. Plate 57, I, shows 16 such units which apparently are equivalent to distinct chromosomes. They separate into two groups, and the elements of the groups migrate along the spindle to the corresponding poles (Pl. 57, J, K). Radiations from the region of the centrosomes are present, but not easily stained during metaphase (Pl. 57, I), and quite long and prominent throughout anaphase (Pl. 57, K) and at telophase (Pl. 57, L, M). They seem to be due to cytoplasmic condensation rather than of the nature of true astral rays. The two chromatin groups in late anaphase are condensed into irregularly lobed masses (Pl. 57, L, M) at the ends of a more or less curved strand of suspension fibers, recalling Blackman's figure 31, Plate 21, (2). The suspension fibers disappear, the daughter nuclei reorganize around the chromatin groups as centers, and a wall divides the young promycelium into two cells (Pl. 57, N). A second division follows immediately (Pl. 57, O, P, Q). The two pairs of granddaughter nuclei thus formed are then separated by cross walls (Pl. 57, R), and the 4-celled promycelium completed. As a rule, the empty basal portion of the promycelium is also cut off by a wall, so that the promycelium really has five cells, four active and one, the stalk, practically dead. Each nucleus is relatively small, with a definite nucleolus and a minute centrosome. The contents take the form of a fine granular network.

From each of the four active cells of the promycelium a stout sterigma is protruded (Pl. 57, S), on which a single spherical sporidium is formed (Pl. 57, V). The nucleus of the cell migrates through the sterigma (Pl. 57, U), taking on an irregular shape during the process, and then rounds up into its normal form (Pl. 57, AA). The division of this nucleus to form a binucleate sporidium (Pl. 57, GG) is quite common, and the karyokinetic figures are particularly striking. Early and late anaphase stages are represented in figures EE and FF, Plate 57. The nuclear behavior in the formation of secondary sporidia was not followed. In germinating sporidia the nucleus probably migrates into the germ tube and there divides, although the figures of this process were indefinite and unsatisfactory. Figure BB of Plate 57 illustrates a common condition which indicates that the nucleus is preparing to divide, at the same

time moving toward the germ tube. At present no data can be given on the young mycelium from the uninucleate or binucleate sporidia, and this comparison must remain impractical until the process of sporidial germination and infection of the pine can be more closely followed.

NUCLEAR PHENOMENA IN VEGETATIVE MYCELIUM IN *PINUS STROBUS*

The uninucleate mycelium produced in the bark of *Pinus strobus* following infection by the sporidia has already been described. The nuclei are completely organized, with a definite nucleolus, chromatin network, and membrane (Pl. 57, HH, II). The chromatin appears to be more or less definitely centered on one point on the nuclear membrane where the centrosome is located. This condition of the arrangement of the chromatin elements of the nucleus will be referred to as polarization, and will be discussed in connection with its appearance in other stages in the cytological cycle. Few cases of vegetative division have been observed in the vegetative cells, but the stages of the process which have been seen indicate that there is a typical rust spindle and that the division is comparable to the vegetative divisions described by Olive (41).

NUCLEAR PHENOMENA IN PYCNIDIUM

The nuclei in the pseudoparenchyma layer from which the short branches bearing the pycnial sporophores arise are quite similar to those of the vegetative mycelium in shape, size, and organization, but they are colored more intensely with hematoxylin and the violet of Flemming's triple stain. The nucleus of the pycnial sporophore is relatively large for the size of the sporophore (Pl. 58, A). In its resting state the chromatin is scattered throughout the whole nucleus in minute granules, which are rather difficult to stain clearly. Each nucleus has a definite centrosome indicated by the local condensation of the chromatin (Pl. 58, A, a). The pycniospore (Pl. 58, A, b), which swells to its full size on the tip of the sporophore, receives one of the daughter nuclei resulting from the division of the sporophore nucleus (Pl. 58, A, d). The exact details of the process were not definitely followed, on account of the small size of the pycniospores and the narrow passage from sporophore to spore. The mature subpyriform pycniospore has the typical structure of corresponding forms in other genera of the rusts. A relatively large nucleus is surrounded by a small amount of cytoplasm (Pl. 58, A, b).

NUCLEAR PHENOMENA IN ÆCIUM

In the layer of fertile cells and in the mycelium below this layer the nuclei are somewhat larger (Pl. 58, H) and more readily stainable than those of the deeper vegetative mycelium (Pl. 57, HH). All the nuclei in this region of the layer and below it exhibit the phenomena of polarization, although the fact may not be evident unless the profile view can be seen.* The same condition persists in all other nuclei in or near the vari-

ous sori. Above the fertile layer a series of cells, the sterile cells (Pl. 58, G), arranged in more or less definite vertical rows, are cut off (Cf. Pl. 54, B, sc). Fromme (14) finds that the sterile cells in *Melampsora lini* (DC.) Tul. arise from the division of the first sterile cells; Moreau (40) states that in *Phragmidium subcorticium* they are formed by the division of the cells in the fertile layers. In *Cronartium ribicola* they seem to arise from the cells of the fertile layer (Pl. 58, F), but there is some evidence that the sterile cells themselves also divide occasionally. Each row contains from 6 to 12 cells, whose contents degenerate as the aëciospores mature. The pressure of the peridium and the lengthening aëciospore chains flatten the whole tissue of sterile cells against the overlying cortex cells.

CELL FUSION TO FORM BASAL CELLS

Fusion normally occurs between two adjacent cells of the fertile layer, which may or may not be at the same level (Pl. 58, I, J). The walls between the fusing cells appear to dissolve, leaving one large cell where there had been two. The basal cell thus formed is the initial cell of the dikaryon. The nuclei are comparatively large (cf. fig. G and I, Pl. 58). They exhibit the polarization phenomena referred to above, but their content is less easily stainable than in the vegetative nuclei below the fertile layer (cf. fig. H and I, Pl. 58), for at times the space between the nucleolus and the membrane seems almost empty (Pl. 58, Y, a). When the nuclei are located in dense cytoplasm toward the upper end of the basal cell, the lower end may be occupied by one or more large vacuoles (Pl. 58, Q). When fusion of more than two cells occurs, the nuclei usually are more irregularly placed (Pl. 58, K, L). The nuclei in these compound basal cells and in the multiple fusion cells are similar to those in the normal binucleate cell.

THE CONJUGATE DIVISION IN THE BASAL CELLS

The division of one of the nuclei of the basal cell will be described, it being understood that the companion nucleus undergoes the same changes at the same time; that is, that the division process is a conjugate division typical of the rusts. On the nuclear membrane is a body which stains deeply with hematoxylin—the centrosome (Pl. 58, I). From it in most cases there run strands of chromatin which are more dense than the other chromatin elements of the nucleus. The nucleus increases in size and the membrane bulges, except at the points where the chromatin strands touch it (Pl. 58, Y, a). At this stage the membrane becomes very thin and then disappears, fading first on the side away from the centrosome (Pl. 58, M, a). Meanwhile the chromatin elements group around or at one side of the nucleolus. Fine suspensors appear to connect the centrosome and the condensing chromatin mass. In the left nucleus in Pl. 58, M, a, the centrosome appears to have divided into two. Coincident with

the disappearance of the nuclear membrane the nucleolus is cast out (Pl. 58, N), after which it migrates off to some distance from the chromatin mass. A true spindle is then formed at the two ends of which are located the centrosomes (Pl. 58, O, P). The original centrosome was not seen to divide, but the conformation of the spindle in some early stages suggests the division of the centrosome to form two, which move apart and become oriented as the opposite poles of the spindle. The chromatin now appears in deeply stained sections as a convoluted dark mass at the middle of the spindle (Pl. 58, O). With better differentiation this mass resolves itself into a jumble of more or less rounded units, the chromosomes (Pl. 58, P). This stage corresponds to the equatorial plate stage of the metaphase. The chromosomes are now seen to separate and move toward the poles (Pl. 58, Q-S); they seem to flow along the outer surface of the spindle rather than to be drawn definitely apart by attraction fibers. Unless very carefully differentiated, the chromatin at this stage may appear to be a single knotted thread reaching from pole to pole. The chromosomes, however, do not fuse into one mass, but, as can be seen in favorable preparations, remain as separate units (Pl. 58, R). As they approach the poles the chromosomes apparently arrange themselves into two groups (Pl. 58, T) for each pole. The division of the centrosomes before the daughter nuclei are reorganized may explain the locus of the two groups, as suggested by Olive (41). At this late stage of anaphase the whole figure may resemble the silhouette dumbbell figures by earlier investigators of the subject. The two groups at each pole now become condensed to two deeply staining masses (Pl. 58, U). Fibrous connections between the daughter groups may still remain visible. The cast-out nucleolus has persisted in the cytoplasm up to this time, slowly becoming less dense and often decreasing in size, while the process described above has been taking place, but as the last spindle fibers disappear and the daughter nuclei begin to become reorganized it fades away, completely absorbed by the cytoplasm. The companion nucleus of the dikaryon, having divided simultaneously with the one described, has by this time given rise to two other daughter nuclei. The four reorganizing nuclei now move apart, two and two, the sister nuclei separating and moving in opposite directions. A wall now forms (Pl. 58, V, a), separating the upper pair from the lower, forming the aëciospore initial and, below it, a new basal cell (Pl. 58, M), potentially equivalent to the primary basal cell.

STEPS IN FORMATION OF AECIOSPORE CHAIN

Division of the dikaryon in the aëciospore initial follows the same process as that in the basal cell (Pl. 58, W, M, b, X). The wall which forms between the two pairs of nuclei divides the original initial cell into two parts, an upper cell which is the aëciospore and a lower, which is the intercalary cell (Pl. 58, Y, d, c). Repeated division of the basal cell and

æciospore initials gives rise to a row of alternating æciospores and intercalary cells (Pl. 58, V), as has been stated in the description of æciospore formation elsewhere in this paper. The nuclei of the æciospores soon become completely reorganized (Pl. 59, AA) and typical of the dikaryon in all its stages, but the nuclei of the intercalary cells slowly degenerate and disappear. A centrosome, the center for the polarization phenomena previously mentioned, can be differentiated by proper staining on the membrane of each nucleus.

NUCLEAR PHENOMENA IN UREDINIUM

The mycelium in leaves of species of *Ribes* has already been described. The nuclei at this stage of the dikaryon are only slightly less in diameter than that of the hyphae, and they are therefore somewhat separated—not side by side as in the elements of the æciospore chain (Pl. 59, C). They do approach each other in the larger cells at the base of the uredinium and undergo conjugate division by a process apparently identical (Pl. 59, E-I) to that described for the basal cell and æciospore initial of the æcium. The same holds true for the division in the urediniospore initial (Pl. 59, J-M, N, b). The wall formed between the two pairs of daughter nuclei divides the initial into an upper, larger cell, the young urediniospore, and a lower, smaller cell, the young stalk cell, which rapidly elongates. The nuclei in the spore become organized similarly to those of the æciospore (Pl. 58, AA), while the stalk nuclei slowly degenerate (Pl. 59, N). The second and subsequent conjugate divisions in the basal cell (Pl. 59, O) are similar to the primary division, but the nuclei often are not so definitely placed side by side, an irregularity probably due to the tendency of the cytoplasm flowing into the second urediniospore initial to pull one of the nuclei along with it. As a rule, both in the basal cell of the aëial chain and in the basal cell of the uredinium at the time of the primary division, the nuclei, though they may commence to divide when located at different levels in the cell, become arranged side by side at metaphase. In the second division of the urediniospore basal cell this orientation may not take place, so that at telephase one of the daughter nuclei may be well up in the new urediniospore initial and its companion just passing into it; but on reorganization after the initial is cut off from the basal cell, the two nuclei take a position side by side in the typical manner. Their division and the subsequent cutting off of the stalk cell by a wall give rise to a second urediniospore. This process may be repeated several times, for the basal cell seems to retain its powers of division until the sorus dries up.

NUCLEAR PHENOMENA IN TELIUM.

The telia are borne on the same mycelium that gives rise to uredinia. By an exactly similar process of conjugate division (Pl. 59, S-W) a cell is cut off from the basal cell of one of the telial unit columns; but this

cell does not divide again, being in itself the young teliospore (Pl. 59, X). The two nuclei appear to become fully reorganized (Pl. 59, Y). They then fuse immediately to produce the single large nucleus of the mature teliospore, during which process the nuclear membranes are absorbed at the points of contact, so that the nuclear contents are free to mix (Pl. 59, Z). The fusing contents round up and become surrounded by the reorganizing membrane. In Plate 59, Z, the two centrosomes and the two nucleoli are still visible. The process of the fusion of the latter was not observed. The fusion nucleus increases in size in a remarkable manner immediately after the union of the two contributing nuclei; and the enlargement may continue until the volume of the fusion nucleus is as much as four times the combined volumes of the two contributing nuclei. Its diameter at the time of maximum enlargement nearly equals the diameter of the teliospore. The chromatin at this stage stains very heavily and appears to be condensed into a heavy irregular spireme-like structure, in which the individual strands are frequently knotted (Pl. 59, AA). Holden and Harper (21), in discussing the fusion nucleus in the teliospore of *Coleosporium sonchi-arvensis* Lev. (*C. solidaginis* (Schw.) Thum.?), present evidence that the heavy skein breaks up into long pieces, which later split longitudinally into finer threadlike units. In *Cronartium ribicola* there certainly occurs a marked reduction in the thickness of the chromatin threads, but no process which could be interpreted as actual splitting was clearly seen. As the chromatin becomes more finely drawn it contracts into a more compact tangle, the nucleus shrinking meanwhile (Pl. 59, BB), and finally splits into granules (Pl. 59, CC). The nucleus at this stage is nearly spherical; the nucleolus is rather small, and is generally located near the membrane. The centrosome could not be differentiated in the fusion nucleus with any stain. So far as could be determined by careful examination, it does not reappear until the primary division in the promycelium and in the resting nuclei of the promycelium.

DISCUSSION OF CYTOLOGICAL PHENOMENA

The nuclear phenomena accompanying cell fusion and spore production in *Cronartium ribicola* clearly confirm the views generally held in regard to rust cytology. Fusion of the gamete cells in pairs in the bark-inhabiting æcium parallels closely similar phenomena in other types of æcia and analogous sori, while multicellular fusion is perhaps much more common than in other forms investigated. Fromme (14) has called attention to triple-cell fusions and the fact that cells below the fertile layer often contribute to the multinucleate fusion cells of *Melampsora lini*. The behavior of the fusion cells in *C. ribicola* shows that cells below those of the fertile layer are potential gametes. It is quite certain that the multiple fusions observed are regular occurrences in æcia of all sizes and shapes, whether on roots or stems. Dittschlag (11) and Hoffman

(20) both figure two aëciospore chains rising from a large basal cell in *Puccinia falcaria* and *Endophyllum semperivi*, respectively, and there is some evidence that aëciospore chains often arise in like manner in *C. ribicola*. It is hardly possible, however, to compare the multiple fusion cells to the central placental cells reported by Richards (49). Inasmuch as the elements of the aëciospore chains are generally binucleate, the number of polynucleate spores being relatively small, it must be concluded that either the extra nuclei so common in the basal cells degenerate or the complex basal cell gives off more than one binucleate spore chain.

The constant similarity in the process of conjugate division in the basal cells of the aëcium, uredinium, and telium suggests a definite stability of the nuclei of the dikaryon throughout its existence. The presence of the centrosome in the resting and dividing nuclei adds confirmation to the reports of this structure in rust nuclei as given by other writers. While it is impossible to make a definite statement from actual observation of the process, it seems perfectly evident that the centrosomes at each pole may divide, thus forming two loci for the chromatin groups approaching the poles—cf. Olive's figures 5, a, and 10, Plate 22 (41). The theory that these two groups of chromatin represent two simple chromosomes, the interpretation of the phenomenon apparently accepted by Mme. Moreau, does not appear tenable, unless it is assumed that these chromosomes are compound and that they break up into their components at metaphase and reunite at telophase. The observations of Holden and Harper (21), Blackman (2), Christman (4), and Olive (41) and the evidence presented in this paper establish the presence of more than two chromosomes or chromosome equivalents, possibly eight being the haploid number. Moreau's (37) figures indicate that her preparations were apparently not sufficiently differentiated to show all the details in the chromatin masses.

The polarization of the nuclei, so evident in nearly all of the resting stages, recalls the condition illustrated in Olive's figure 8, Plate 22 (41). The arrangement of the chromatin with a point on the membrane as a locus for the convergence of the strands is not due to fixation; for it is a constant phenomenon in nuclei killed and fixed under different conditions and with different reagents. The centrosome is apparently the center of attraction. Harper (28) has conclusively demonstrated similar phenomena in the Erysipheae. He has also called attention to the fact that the nuclei of these ascomycetes exhibit a definite regularity and stability with reference to the arrangement of the chromatin strands and the polarization throughout the cytological changes which he observed. He believes the chromosomes retain their individuality to a marked degree in all the nuclear processes. The observations on the nuclear structure in *Cronartium ribicola* herein presented suggests that

perhaps the conditions in the rust nucleus approach more nearly than has been supposed the conditions in the Ascomycetes and other fungi. The process of nuclear fusion in the young teliospore is comparable to the same process in other rusts. It seems reasonable to conclude that this fusion is the completion of the process, initiated at the time of cell fusion to form the dikaryon, which takes the place of normal fertilization, and that the actual fusing is a necessary preliminary to the mixing of the chromatin elements and the subsequent reduction division, as suggested by Maire (31) and others. Certainly the changes which take place in the nucleus after fusion suggest a complicated mingling and readjustment of the chromatin which would seem to justify such a view. It is not necessary to regard the fusion as a pseudo fertilization (10). There appears to be little reason for doubting that the first division in the promycelium is heterotypical (38), for it is unique and decidedly different from the second division which immediately follows. Arnaud (1) has compared the mitotic figures in *Capnodium meridionale* with those in *Coleosporium senecionis*. Wager's figure 84, Plate 19 (63), of the telophase in the dividing fusion nucleus in the sporangium of *Polyphagus euglenae* is very like similar stages in the primary division in the promycelium of *Cronartium ribicola*. It may be noted that the association of the nuclei in *P. euglenae* and their subsequent fusion in the sporangium are phenomena comparable to the nuclear conditions in the rust dikaryon and fusion in the teliospore.

ABNORMALITIES

Before coming to the general discussion and summary it is necessary to mention briefly certain abnormalities commonly met in the different types of sori. Aëcia sometimes occur with reversed polarity in part of the sorus, where the spores are produced on aëcial chains which grow toward the center of the tree, usually into a resin duct. This change in the direction of growth is probably to be explained by the fact that the developing chains followed the line of least resistance, in this case into the adjacent resin duct, instead of against the overlying host cells. Aëcia also often develop on the roots, under several inches of leaves and loam. Their structure appears to be normal, but their environment is hardly advantageous. Double pycnial layers (Pl. 50, C, a, a₁) are not uncommon. In a short note, Posey, Gravatt, and Colley (45) have reported the finding of uredinia on the stems of *Ribes hirtellum* Michx. In the cortex of infected stems of this species internal uredinia (Pl. 51, C) with normal and reversed polarity were formed in abundance. Internal telia, produced in the pith and cortex of the petioles of *Ribes* sp.,¹ have been described in a previous paper (8). In such abnormal

¹ *Ribes roezli* was the name given in the original article, but the species determination was probably incorrect. As the species has not fruited, accurate determination has been impossible.

sori, typical column development is sometimes completely inhibited by the pressure of the surrounding host tissue. All internal sori are to be regarded as teratological phenomena of no special morphological significance. Uredinia and telia on the petiole are common; occasionally they develop on the upper surfaces of infected leaves.

GENERAL DISCUSSION

The writer's observations on the mycelium of *Cronartium ribicola* in the bark of *Pinus strobus* confirm the conclusions of Klebahn (25) in regard to the course of the hyphæ of the parasite and add considerable new data on the inter relations of the host and parasite cells. Hartig's paper (19) on *Peridermium pini* and Wolff's article (64) on the same fungus, which appeared shortly after Hartig's and in some places is almost a direct copy of Hartig, are the only papers which have come to the writer's attention which describe and illustrate the morphology and parasitic relations of a bark-inhabiting rust and its pine host. Hartig believes that the swelling of the bark of the host is due to the fact that the cortex and phloem cells are forced apart by the abundant intercellular mycelium of the parasite. This has been shown to be the case with the hypertrophy produced by *C. ribicola* in the bark of *P. strobus*. Observations on the bark of *P. parviflora* Sieb. and Zucc. infected with the same fungus show that the swelling is produced in the same way as it is in *P. strobus*. The same statement holds true for the swelling caused by *C. complanata* Arth. in the bark of *P. sylvestris* L., and *P. ponderosa* Laws. Neither *C. ribicola* nor *C. complanata* are gall-forming rusts. Their mycelium is confined almost entirely to the region of the cortex and phloem cells of their pine hosts, although the hyphæ do enter the wood along the rays, as Tubeuf (58) has reported for *Peridermium pini*, and occasionally work their way in between tracheids. These hyphæ have been traced in the case of *C. ribicola* to a depth of three annual rings, counting in from the cambium. In all probability it will be found that the mycelium may be found in the ray cells of the annual ring laid down at about the time infection took place, although it may not remain active after the ray cells die. While the presence of the hyphæ in a given annual ring may not mean that the pine was infected during the year this annual ring was laid down, because there must be some growth along the ray cells toward the center of the tree, it will establish roughly the date of infection. The similarity which evidently exists between the morphology and method of parasitism of *C. ribicola* and *C. complanata* suggests that a close agreement will probably be found among all non-gall-forming caulicolous *Peridermia*.

The actual injury to the pine host cells from the irritation caused by the invading mycelium is apparently very slight. The cells pierced by

haustoria continue to remain alive and apparently active in spite of haustoria. As has been pointed out above, these haustoria become invested with a sheath which may render them very ineffective absorbing organs, in which case they might become simple mechanical irritants in the cell content, no more detrimental to cell activity than crystals. The denting of the host nucleus by the tips of the haustoria does not seem to injure the nucleus except to alter its shape. There is no evidence of increased cell division reported by Reynolds (48) for many plants as a direct result of a parasite's action, or of nuclear migrations like those figured by Schürhoff (52). Storage starch is usually present in excess, but this phenomenon has been shown by Halsted (17) to be a general condition in and around areas infected with fungus parasites. This starch is not completely used up by the fungus, for many grains remain in the old dead cells after the cells are completely dried out. The cells just beneath the pycnial layer and in the region of the young æcium (Pl. 52, B) contain normal grains as well as decomposition products. Excess starch production is probably due to a lack of balance in the physiological processes in the host cells, and the fungus may, of course, contribute to the unbalancing; but other environmental factors which are little understood result in excess starch production in trees which appear to be perfectly normal in other respects.

The wood laid down in the annual rings under infected bark is much less than in healthy trees, but the tracheids are apparently normal in everything except number. The presence of the hyphæ in the tracheids has no appreciable influence on their form; neither are the characters of the ray cells perceptibly changed.

Cronartium ribicola may be the primary cause of the death of a young tree. However, the swelling of the bark is not in itself a serious hindrance to conduction in the phloem. The actual severe injury occurs when the æcia form and burst through the outer bark; for the æcial cracks thus formed allow the inner bark cells to dry out and die. This results in the breaking of resin canals and the consequent exudation of resin in large quantities. The girdling of the tree is due to these two causes working together—namely, the cracking and drying of the bark, and the impregnation of the whole cortex and phloem in the cracked area with resin. Complete stoppage of the conducting elements of the phloem results. Seedlings and young trees may succumb to the attack of the fungus almost as soon as the first æcia appear; but with older trees death is sometimes delayed for a number of years. It depends, of course, on how complete a girdle has been effected. The part played by secondary fungi acting in conjunction with *C. ribicola* to make a girdle complete is very important. It will be recognized at once that the cracking of the bark at the time the æcia are formed is a source of danger to the trees not only by exposing the inner bark cells to the air, but also by providing an avenue of entrance for secondary

parasitic fungi, or saprophytic fungi, and insects, which hasten the decomposition of the weakened susceptible tissue. These secondary fungi may also gain entrance through the pycnial spots. Rathay (47) has shown that the pycnia of many rusts are visited by insects, apparently attracted by the sweet drops. Both pycnia and æcia in *C. ribicola* may be eaten out by various insects, and unquestionably these insects bear on their bodies or legs viable spores of fungi capable of growing in the bark tissue. In many instances these fungi are so rapid in their growth that they overrun the infected area and completely suppress the rust, so that it never forms æcia. In such cases, the bark shrinks and the infected stem is actually constricted, and the girdle thus formed is often more quickly effective than in the cases where rust works alone.

In leaves of *Ribes* spp. the mycelium sometimes causes the death of isolated infected spots; but in other cases the hyphæ penetrate to all parts of the tissue without causing death of the cells, and without producing hypertrophy. The large spaces among the mesophyll cells and the fact that the hyphæ rarely form solid mycelial masses in leaves of *Ribes* spp. probably help to explain the lack of hypertrophy and destructive effect. When defoliation occurs during the course of severe epidemics of the rust, there is, of course, a consequent poor crop of berries. The variation in the effect of the parasite on the different species of *Ribes* is a subject which must have separate treatment and therefore can not be considered fully at this time.

It will probably be found on further investigation that a close agreement exists among caulicolous *Peridermia* with respect to the structure of the pycnium. The external appearance of the pycnium, or of the thin layer of tissue overlying it, will, however, be found to vary according to the outer bark texture of the host. In the case of *Pinus parviflora* the exterior appearance of the pycnium is almost identical with the pycnium of *P. strobus*. The value of the pycnial spots as diagnostic characters has been briefly outlined in an earlier short note (6), and referred to above in connection with the discussion of the formation of the pycnium and pycniospores. Hartig (19, *Taf.* 4, *fig.* 7, b) was one of the first observers to call attention to the pycnial spots on the bark of *P. strobus*, although he at that time thought the fungus on this pine was identical with *Peridermium pini*. Wolff (64) copied the same figure in his paper on the latter fungus. Kirchner and Boltshauser (22) in Plate 15 of their atlas show what are evidently three pycnial spots, but they do not definitely refer to them in their description of the figures. The ability to recognize the pycnial spots of *Cronartium ribicola* is almost absolutely essential in the field study of control methods.

The description of the formation of the æcium given above emphasizes the remarkable agreement which exists in the fundamental processes involved in the production of æciospores in the rusts. The figures of Hartig (19), Wolff (64), and Sappin-Trouffy (51, *fig.* 65) on the æcium of

Peridermium pini are not complete nor detailed enough to enable one to compare directly the structure of the acicolous and caulicolous æcia with the structure of the æcium of *Cronartium ribicola*. It is evident, however, that the leaf and stem types are quite similar. Study of the expansive æcia of *C. occidentale* Hedge., Bethel, and Hunt may reveal some interesting morphological variations.

The details given for the formation of the multilayered peridium are apparently the first published record of the origin of this structure. Emphasis should perhaps also be placed on the constant and normal occurrence of multinucleate cells at the base of the æcium, a phenomenon which has been discussed elsewhere, by suggesting that in deep-seated æcia of caulicolous *Peridermia* the æciospore chains may be found to arise more often from these placenta-like cells than from basal cells arising as a result of the fusion of only two cells in the fertile layer. Although recent investigators—for example, Kurssanow (27)—have confirmed the results of Christman (4) and other writers with regard to the origin of the basal cell as a result of the fusion of two adjacent fertile cells, it yet remains a question whether this method is constant or whether certain variations in the formation of the basal cells are to be expected in deep-seated æcia.

Ludwig and Rees (28) in a recent article report some details of the structure of the uredinium of *Pucciniastrum agrimoniae* (Schw.) Tranz. Their figure would serve very well for a figure of the young uredinium of *Cronartium ribicola* (cf. Pl. 55, B). In the description of the uredinium of the latter it has been shown that the peridium is formed by the coalescence of cells which are cut off from certain cells that are analagous to the basal cells of the sorus. These cells also cut off urediniospore initials which then divide into urediniospores and stalk cells. In the young sorus these four cells—that is, the basal cell, the stalk cell, the urediniospore, and the peridial cell—form what looks like a chain of cells. As the spores and their stalks mature, the row arrangement is lost in the middle of the sorus, but persists at the circumference. This fact places the conclusion of Ludwig and Rees that the urediniospores of *P. agrimoniae* are borne in chains, under suspicion. Personal investigation by the writer into the structure of the uredinium in this species of *Pucciniastrum* shows that the method of formation of the urediniospores in *P. agrimoniae* and in *C. ribicola* is practically identical, and that therefore the spores in the uredinium of the former are not borne in chains but on stalks. In the case of *P. agrimoniae* the stalks are sometimes quite short and the basal cells from which they arise are much less conspicuous than they are in *C. ribicola*, but these differences are not important as far as the method of spore formation is concerned. The encircling bank of parenchyma-like cells surrounding the uredinium in *C. ribicola* is not found in *P. agrimoniae*. It is difficult to interpret Magnus's (30) figures of the uredinia of the *Pucciniastrum* group as to the exact morphology of the cells from

which the urediniospores arise, but his description of the manner in which the urediniospores in Pucciniastrum are borne is correct. Fischer's (13) figure of the uredinial peridium of *C. asclepiadeum* (Willd.) Fr. evidently represents the same type of peridium as is present in the uredinium of *C. ribicola*. The peridial cells surrounding the break in the top of the peridium in the latter fungus are thickened irregularly, but they do not often appear as conspicuous as the large cells figured by Fischer (13) in the case of the uredinial peridium in species of Pucciniastrum or by Ludwig and Rees (28) for *P. agrimoniae*.

In the discussion of the telium it has been shown that it originates in a way which makes it impossible to tell the young telium from the young uredinium. Under these circumstances it is most natural that a peridium should be found over the telium, as in the case of the young uredinium. The same holds true for the parenchyma-like bank of cells which encircle the base of the column. Both the peridium and the bank of encircling cells come into being before the sorus becomes differentiated into either uredinium or telium. With these facts clearly in view it may be reasonably safe to predict that a telial peridium, a structure which has apparently not been reported previously, will be found to be present in other species of the genus *Cronartium*. The manner of the germination of the teliospores, previously discussed, adds to our knowledge of the morphology and behavior of the spores in the telial column, at the same time confirming the observations of Tulasne (61) and Sappin-Trouffy (57) on other species.

This paper is offered as a contribution to our knowledge of the parasitism, morphology, and cytology of the rusts, and especially of the genus *Cronartium*. It is hoped that the data presented may prove valuable in stimulating further research on the interrelations of rust parasites and their hosts, which will, of course, involve more accurate study of the anatomy of the hosts and the modifications in the normal structure of the host tissues under the action of the parasites. The need for a comparative paper on the haustoria of the rusts scarcely requires emphasis. It will be interesting to compare the results of investigations on the cytology of other deep seated cancolous æcia with those herein presented, especially with reference to the process of the formation of basal cells, the phenomena of polarization of the nuclei, and the centrosomes and chromosomes.

SUMMARY

(1) In the foregoing paper hitherto unpublished data on the morphology and cytology of *Cronartium ribicola* Fischer and the interrelations of the parasite and its hosts, *Pinus strobus* and *Ribes* spp., are presented and fully illustrated.

(2) The mycelium is more abundant in the *Pinus strobus* than in species of *Ribes*. In the former the hypæ force the cortex and phloem

cells apart and thus cause the swelling of the infected bark; in the latter there is rarely any marked aggregation of the hyphae, except in the case of petiolar infection.

(3) Haustoria may penetrate practically every cell in the infected area in *Pinus strobus*. These haustoria are characteristic for *Cronartium ribicola* and their presence in the bark cells of the pine definitely determines the identity of the parasite. A sheath develops around each haustorium as it reaches maturity or old age. This sheath is apparently unlike any other sheath so far described in connection with the haustoria of the rusts.

(4) Haustoria are proportionately much less frequent in *Ribes* spp. than in *Pinus strobus* and much smaller in size. They are not enveloped in sheaths as in the latter.

(5) The morphology of the different sori is shown to be similar to the morphology in other full-cycle rusts.

(6) The structure of the spreading pycnial layer characteristic of caulicolous *Peridermia* is considered in detail.

(7) The development of the deep-seated aecium and the formation of its multilayered peridium is described.

(8) The formation of urediniospores is shown to follow the general plan in other uredinia where the spores are borne on stalks. The development of the uredinial peridium and the bank of parenchyma-like cells which surround the uredinium are described and figured for the first time.

(9) The telial column is shown to arise either from an old uredinium or as a separate entity and to be indistinguishable from a young uredinium in its very early stages. It is surrounded at the base with a bank of parenchyma-like cells and is provided with a peridium similar to the peridium of the uredinium. All of the spores of the telial column may germinate *in situ*. The production of sporidia is described in detail.

(10) The destructive effect on the pine host as a result of the attack of *Cronartium ribicola* varies. In young trees death may result quickly. In older trees it is of the nature of a primary injury, which prepares the way for the drying out of the infected bark and the entrance of secondary fungi and insects which complete the destruction initiated by the parasite.

(11) The effect on *Ribes* varies with the species attacked. It may result in early defoliation and a consequent poor crop, but in general it is not serious on this host.

(12) The cytological phenomena presented herein establish the similarity of the nuclear processes in the genus *Cronartium* and other full-cycle rusts.

(13) The resting nuclei in all cases, with the exception of the mature fusion nucleus in the teliospore, have a deeply staining spot on the

membrane called the centrosome. This body is apparently the center for the more or less definite convergence and concentration of the chromatin strands referred to as polarization.

(14) The presence of large, irregular, multinucleate cells at the base of the æcium is a regular phenomenon and not to be regarded as an abnormal condition. Probably more than one æciospore chain may arise from such cells.

(15) There is a remarkable constancy in the process of conjugate division and stability in the nuclear structure throughout the dikaryon. In the dividing nuclei the centrosomes are visible as deeply staining dots at the poles of the spindle. The number of chromosomes is certainly more than two. Possibly eight is the haploid number.

LITERATURE CITED

- (1) ARNAUD, G.
1913. LA MITOSE CHEZ CAPNODIUM MERIDIONALE ET CHEZ COLEOSPORIUM SENECTIONIS. *In* Bul. Soc. Mycol. France, t. 29, fasc. 3, p. 345-347, pl. 22-23.
- (2) BLACKMAN, V. H.
1904. ON THE FERTILIZATION, ALTERNATION OF GENERATIONS, AND GENERAL CYTOLOGY OF THE UREDINEAE. *In* Ann. Bot., v. 18, no. 71, p. 323-373, pl. 21-24. List of papers, p. 368-369.
- (3) ——— and FRASER, Helen C. I.
1906. FURTHER STUDIES ON THE SEXUALITY OF THE UREDINEAE. *In* Ann. Bot., v. 20, no. 77, p. 35-48, pl. 3-4.
- (4) CHRISTMAN, A. H.
1905. SEXUAL REPRODUCTION IN THE RUSTS. *In* Bot. Gaz., v. 39, no. 4, p. 267-275, pl. 8.
- (5) ———
1907. THE ALTERNATION OF GENERATIONS AND THE MORPHOLOGY OF THE SPORE FORMS IN THE RUSTS. *In* Bot. Gaz., v. 34, no. 2, p. 81-101, pl. 7. Literature cited, p. 99-100.
- (6) ———
1907. THE NATURE AND DEVELOPMENT OF THE PRIMARY UREDO-SPORE. *In* Trans. Wis. Acad. Sci., v. 15, pt. 2, p. 517-526, pl. 29.
- (7) COLLEY, Reginald H.
1917. DIAGNOSING WHITE-PINE BLISTER-RUST FROM ITS MYCELIUM. *In* Jour. Agr. Research, v. 11, no. 6, p. 281-286, 1 fig., pl. 31.
- (8) ———
1917. DISCOVERY OF INTERNAL TELIA PRODUCED BY A SPECIES OF CRONARTIUM. *In* Jour. Agr. Research, v. 8, no. 9, p. 329-332, pl. 88. Literature cited, p. 332.
- (9) ———
1917. PYCNIAL SCARS, AN IMPORTANT DIAGNOSTIC CHARACTER FOR THE WHITE PINE BLISTER RUST. (Abstract.) *In* Phytopathology, v. 7, no. 1, p. 77.
- (10) DANGEARD, P.-A., and SAPPIN-THOUFFY.
1893. UNE PSEUDO-FÉCONDATION CHEZ LES URÉDINÉES. *In* Compt. Rend. Acad. Sci. [Paris], t. 116, no. 6, p. 267-269.
- (11) DITTSCHLAG, E.
1910. ZUR KENNNTNIS DER KERNVERHÄLTNISSE VON PUCCINIA FALCARLÆ. *In* Centbl. Bakt. [etc.], Abt. 2, Bd. 28, No. 16/19, p. 473-492, 7 fig., pl. 1-3.

- (12) DUFF, George H.
1918. SOME FACTORS AFFECTING VIABILITY OF THE UREDINIOSPORES OF CRO-
NARTIUM RIBICOLA. In *Phytopathology*, v. 8, no. 6, p. 289-292, 1 fig.
- (13) FISCHER, Ed.
1904. DIE UREDINEEN DER SCHWEIZ. . . 590 p., illus. Bern.
- (14) FROMME, Fred D.
1912. SEXUAL FUSIONS AND SPORE DEVELOPMENT OF THE FLAX RUST. In
Bul. Torrey Bot. Club, v. 39, no. 3, p. 113-131, pl. 8-9. Literature
cited, p. 129.
- (15) FROMME, F. D.
1914. THE MORPHOLOGY AND CYTOLOGY OF THE AECIDIUM CUP. In *Bot. Gaz.*,
v. 58, no. 1, p. 1-35, 8 fig., pl. 1-2.
- (16) GUILLIERMOND, A.
1913. LES PROGRÈS DE LA CYTOLOGIE DES CHAMPIGNONS. In *Prog. Rei Bot.*,
Bd. 4, Heft 3/4, p. 389-542, 82 fig. Index bibliographique, p. 534-542.
- (17) HALSTED, Byton D.
1898. STARCH DISTRIBUTION AS AFFECTED BY FUNGI. In *Proc. Amer. Assoc.*
Adv. Sci., 47th Meeting, p. 408-409.
- (18) HARPER, R. A.
1905. SEXUAL REPRODUCTION AND THE ORGANIZATION OF THE NUCLEUS IN
CERTAIN MILDREWS. 104 p., 7 pl. Washington, D. C. (Carnegie Inst.
Washington Pub. 37.)
- (19) HARTIG, Robert.
1874. WICHTIGE KRANKHEITEN DER WALDBÄUME. 127 p., 6 pl. Berlin.
- (20) HOFFMAN, A. W. Hans.
1911. ZUR ENTWICKLUNGSGESCHICHTE VON ENDOPHYLLUM SEMPERVIVI. In
Centbl. Bakt. [etc.], Bd. 32, No. 3/5, p. 137-158, 14 fig., 2 pl. Litera-
turverzeichnis, p. 157-158.
- (21) HOLDEN, R. J., and HARPER, R. A.
1903. NUCLEAR DIVISIONS AND NUCLEAR FUSION IN COLEOSPORIUM SONCH-
ARVENSIS LEV. In *Trans. Wis. Acad. Sci.*, v. 14, pt. 1, 1902, p. 63-82,
pl. 1-2.
- (22) KIRCHNER, O., and BOLTSCHAUER, H.
1902. ATLAS DER KRANKHEITEN UND BESCHÄDIGUNGEN UNSERER LAND-
WIRTSCHAFTLICHEN KULTURPFLANZEN. . . 8. 6. Stuttgart.
- (23) KLEBAHN, H.
1888. BEOBSACHTUNGEN UND STREITFRAGEN ÜBER DIE BLASENROSTE. In
Abhandl. Ver. Bremen, Bd. 10, Heft 1, p. 145-155, pl. 1.
- (24) ———
1902. KULTURVERSUCHE MIT HETEROCISCHEN UREDINEEN. In *Ztschr. Pflan-
zenkrank.*, Bd. 2, p. 258-275, 332-343, pl. 5.
- (25) ———
1904. DIE WIRTSWECHSELNDEN ROSTPILZE. . . xxxvii, 447 p., illus. Berlin.
Literature, p. ix-xxxvii.
- (26) ———
1905. KULTURVERSUCHE MIT ROSTPILZEN. In *Ztschr. Pflanzenkrank.*, Bd. 15,
Heft 2, p. 65-108, 4 fig., pl. 3.
- (27) KURSSANOW, L.
1910. ZUR SEXUALITÄT DER ROSTPILZE. In *Ztschr. Bot.*, Jahrg. 2, Heft 2,
p. 81-93, pl. 1.
- (28) LUDWIG, C. A., and REES, C. C.
1918. THE STRUCTURE OF THE UREDINIUM IN FUCCINIASTRUM AGRIMONIAE.
In *Amer. Jour. Bot.*, v. 5, no. 2, p. 55-60, pl. 8.

- (29) McCUBBIN, W. A.
1917. CONTRIBUTIONS TO OUR KNOWLEDGE OF THE WHITE PINE BLISTER RUST.
In Phytopathology, v. 7, no. 2, p. 95-100, 1 fig.
- (30) MAGNUS, P.
1909. BEMERKUNGEN ÜBER EINIGE GATTUNGEN DER MELAMPFOREEN. *In*
Ber. Deut. Bot. Gesell., Bd. 27, Heft 6, p. 320-327, pl. 14.
- (31) MAIRE, René.
1900. L'ÉVOLUTION NUCLÉAIRE CHEZ LES ENDOPHYLLUM. *In* Jour. Bot.,
ann. 14, no. 3, p. 80-92; no. 4, p. 94-97; no. 12, p. 369-382, pl. 3.
- (32) ———
1911. LA BIOLOGIE DES URÉDINALES. *In* Prog. Rei Bot., Bd. 4, Heft 1, p.
109-162. Index bibliographique, p. 160-162.
- (33) ———
1912. [REVIEW OF PAVILLARD, J. REMARQUES SUR L'ÉVOLUTION DES URÉ-
DINÉES.] *In* Mycol. Centbl., Bd. 1, Heft 7/8, p. 214.
- (34) MAIRE, R.
1913. LA STRUCTURE ET LA POSITION SYSTÉMATIQUE DU MAPEA RADIATA PAT.
In Bul. Soc. Mycol. France, t. 29, fasc. 3, p. 335-338, 1 fig.
- (35) MARCHAL, Émile.
1901. RAPPORT SUR LES MALADIES CRYPTOGAMIQUES ÉTUDIÉES AU LABORATOIRE
DE BOTANIQUE DE L'INSTITUT AGRICOLE DE GEMBOUX. *In* Bul. Agr.
[Brussels], t. 17, p. 4-18, 5 fig.
- (36) MOREAU, Fernand.
1913. LE CENTROsome CHEZ LES URÉDINÉES. *In* Bul. Soc. Mycol. France,
t. 29, fasc. 2, p. 242-243. Index bibliographique, p. 243.
- (37) ———
1913. LES PHÉNOMÈNES DE KARYOKINÈSE CHEZ LES URÉDINÉES. *In* Bul.
Soc. Bot. France, t. 60, no. 2/3, p. 138-141.
- (38) ———
1914. LA MITOSE HÉTÉROTYPIQUE CHEZ LES URÉDINÉES. *In* Bul. Soc. Bot.
France, t. 61, no. 1/3, p. 70-74.
- (39) ———
1914. LA MITOSE HOMÉOTYPIQUE CHEZ LE COLEOSPORIUM SENECTIONIS PERS.
In Bul. Soc. Bot. France, t. 61, no. 1/3, p. 4-5.
- (40) ———
1914. SUR LE PRÉTENDU TRICHOGYNE DES URÉDINÉES. *In* Bul. Soc. Mycol.
France, t. 30, fasc. 3, p. 368-372.
- (41) OLIVE, Edgar W.
1908. SEXUAL CELL FUSIONS AND VEGETATIVE NUCLEAR DIVISIONS IN THE
RUSTS. *In* Ann. Bot., v. 22, no. 86, p. 331-360, pl. 22. Literature,
p. 358.
- (42) ———
1918. THE CYTOLOGICAL STRUCTURE OF BOTRYORHIZA HIPPOCRATEAE. *In* Mem.
Brooklyn Bot. Gard., v. 1, p. 337-341, pl. 8.
- (43) PAVILLARD, J.
1912. REMARQUES SUR L'ÉVOLUTION DES URÉDINÉES. *In* Bul. Soc. Mycol.
France, t. 28, fasc. 1, p. 57-59.
- (44) POIRAULT, G., and RACIBORSKI, M.
1895. SUR LES NOYAUX DES URÉDINÉES. *In* Jour. Bot., ann. 9, no. 17, p. 318-
324, fig. 1-12; no. 18, p. 325-332, fig. 15-19; no. 20, p. 381-388, pl. 6.
- (45) POSEY, G. B., GRAVATT, G. Filippo, and COLLEY, Reginald H.
1917. URÉDINIA OF CRONARTIUM RIBICOLA ON RIBES STEMS. *In* Science, n. s.
v. 46, no. 1187, p. 314-315.

- (46) RAMSBOTTOM, J.
1913. RECENT PUBLISHED RESULTS ON THE CYTOLOGY OF FUNGUS REPRODUCTION (1913). *In* Brit. Mycol. Soc. Trans., v. 4, pt. 2, p. 249-291.
- (47) RÁTHAY, Emeric.
1883. UNTERSUCHUNGEN ÜBER DIE SPERMOGONIEN DER ROSTPILZE. *In* Denkschr. K. Akad. Wiss. [Vienna], Math. Naturw. Cl., Bd. 46, Abt. 2, p. 1-51.
- (48) REYNOLDS, Ernest Shaw.
1912. RELATIONS OF PARASITIC FUNGI TO THEIR HOST PLANTS. I. Studies of parasitized leaf tissue. *In* Bot. Gaz., v. 53, no. 5, p. 365-395, 9 fig. Literature cited, p. 390-395.
- (49) RICHARDS, Herbert Maule.
1896. ON SOME POINTS IN THE DEVELOPMENT OF *ÆCIDIA*. *In* Proc. Amer. Acad. Arts and Sci., v. 31 (n. s. v. 23), 1895/96, p. 253-270, pl.
- (50) SAPPIN-THOUFFY.
1892. LES SUÇOIRS CHEZ LES URÉDINÉES. *In* Botaniste, s. 3, p. 215-219, pl. 18.
- (51) ———
1896. RECHERCHES HISTOLOGIQUES SUR LA FAMILLE DES URÉDINÉES. *In* Botaniste, s. 5, fasc. 2/5, p. 59-244.
- (52) SCHÜRHOFF, P.
1906. DAS VERHALTEN DES KERNES IN WUNDGEWEBE. *In* Beih. Bot. Centbl., Bd. 19, Heft 3, p. 359-382, pl. 9.
- (53) SMITH, Grant.
1900. THE HAUSTORIA OF THE ERYSPHÆE. *In* Bot. Gaz., v. 29, no. 3, p. 153-184, pl. 11-12. Bibliography, p. 181-183.
- (54) SPAULDING, Perley.
1911. THE BLISTER RUST OF WHITE PINE. U. S. Dept. Agr. Bur. Plant Indus. Bul. 206, 88 p., 5 fig., 2 pl. (1 col.). Bibliography, p. 61-78.
- (55) ———
1916. THE WHITE-PINE BLISTER RUST. U. S. Dept. Agr. Farmers' Bul. 742, 15 p., 5 fig., 1 pl. (col.).
- (56) ——— and GRAVATT, G. Flippo.
1917. INOCULATIONS ON RIBES WITH *CRONARTIUM RIBICOLA* Fischer. *In* Science, n. s. v. 46, no. 1184, p. 243-244.
- (57) STEWART, F. C.
1906. AN OUTBREAK OF THE EUROPEAN CURRANT RUST (*CRONARTIUM RIBICOLA* DIETR.). N. Y. State Agr. Exp. Sta. Tech. Bul. 2, p. 61-74, 3 pl. Bibliography, p. 71-74.
- (58) TUBEUF, Karl v.
1895. PFLANZENKRANKHEITEN . . . 599 p., illus. Berlin.
- (59) TUBEUF, Karl von.
1897. UEBER DIE VERBREITUNG VON PFLANZENKRANKHEITEN. I. DIE GEFAHR DER AUSDEHNUNG DER RINDENBLASENROSTES DER WEY-MUTHSKIEFER. *In* Forstl. Naturw. Ztschr., Jahrg. 6, Heft 9, p. 320-324, 4 fig.
- (60) ———
1905. ÜBER DIE BIOLOGIE, PRAKTISCHE BEDEUTUNG UND BEKÄMPFUNG DES WEIMUTSKIEFERN-BLASENROSTES. K. Biol. Anst. Land u. Forstw. Flugbl. 5, Aufl. 2, 4 p., col. pl.
- (61) TULASNE, L. R.
1854. SECOND MÉMOIRE SUR LES URÉDINÉES ET LES USTILAGINÉES. *In* Ann. Sci. Nat. Bot., s. 4, t. 2, p. 77-196. pl. 7-12.

- (62) VUILLEMIN, Paul.
1893. SUR LA FÉCONDATION DES PUCCINIÈRES. In *Compt. Rend. Acad. Sci.*
[Paris], t. 116, no. 25, p. 1464-1467.
- (63) WAGER, Harold.
1913. THE LIFE-HISTORY AND CYTOLOGY OF POLYPHAGUS EUGLENÆ. In *Ann.*
Bot., v. 27, no. 106, p. 173-202, pl. 16-19.
- (64) WOLFF, Reinhold.
[1876.] *ACIDIUM PINI* PERS. UND SEIN ZUSAMMENHANG MIT *COLEOSPORIUM*
SENECIONIS LÉV . . . 28 p., 2 pl. [Riga.] (Festschrift der kaiser-
lichen Akademie der Wissenschaften zu St. Petersburg zu ihrer hun-
dertundfünfzigjährigen Gedenkfeier dargebracht von der Plenarcon-
ferenz der polytechnischen Schule zu Riga.)

PLATE 48

Cronartium ribicola on *Pinus strobus*:

A.—This figure illustrates the etiolated condition of the bark in the case of a comparatively young nodal infection. The node lies between 4- and 5-year-old wood. The infection originated on the opposite side of the stem to that shown in the photograph. Note the irregular margin of the etiolated area. The mycelium, starting in opposite directions from the infection point, has now completely encircled the stem. The junction point of the encircling hyphæ is along the line *ab*. Natural size.

B.—This figure illustrates an internodal infection, somewhat older than that shown in figure A, in which the infection apparently originated at the base of the leaf fascicle (*a*). The darker patches on the pronounced canker area are pycnial spots. *Æcia* are formed under the bark all over the canker and are beginning to break out toward the upper end. Note the characteristic shape of the young canker. Natural size.






PLATE 49

Cronartium ribicola:

A.—The edge of a pycnium in section. Note the position of the pycnium with reference to the overlying periderm and underlying cells of the outer cortex. $\times 250$.

B.—Part of the same section showing the general relation of the elements which go to make up the sorus and their relation to the host cells beneath. *a*, Contributing hyphæ; *b*, pseudoparenchyma layer; *c*, sporophores. Compare Pl. 58, A. $\times 1,050$.

C.—Tangential section in the xylem, showing the cut end of a ray and the manner in which a haustorium (*a*) may rise from the hyphæ in the ray and enter the lumen of the tracheid. $\times 525$.


PLATE 50

Cronartium ribicola:

A.—A figure illustrating the relation of the pycnium (*a*), the heavy black line at the top, and young æcium, to the host tissue. The heavily stained cells (*b*) in the region of the young æcium are the fertile cells. Compare Plate 54, B. $\times 125$.

B.—A section through a mature æcium, taken a little to one side of the break in the bark, to show the æciospore chains (*a*), the multilayered peridium (*b*), and the overlying host tissue. $\times 75$.

C.—A similar section showing a double pycnial layer (*a*, *a*₁), and the location of the cork cambium (*b*) which cuts out the old pycnium. $\times 40$.






PLATE 51

Cronartium ribicola:

A.—A median section through a young uredinium forming in the space beneath a stoma. *a*, Peridial cells; *b*, young urediniospores. The photomicrograph illustrates a stage in the development of the uredinium midway between the stages shown in Plate 51, B, and 55, B. $\times 650$.

B.—A section through the same uredinium as that shown in Plate 55, C, taken to one side of the break in the peridium, toward the edge of the sorus. *a*, Peridial cells; *b*, a young urediniospore; *c*, A stalk cell. $\times 525$.

C.—An internal uredinium from the cortex of *Ribes hirtellum*. $\times 315$.

PLATE 52

Cronartium ribicola:

A.—A section of a young telial column. Note the overlying peridial cells (*a*) and the row arrangement of the developing teliospores. The epidermal cells have been torn off. $\times 355$.

B.—A later stage in the development of the telial column. Note the shape of the tip cells and the character of the parenchyma-like cells (*a*) surrounding the base of the young column (cf. Pl. 55, C). The cells near the base of the column are binucleate, while those at the tip are uninucleate. $\times 355$.

C.—A longitudinal section of a mature column. $\times 250$.

D.—Higher power view of the same section, showing the arrangement of the individual spores, and the size of the nuclei. $\times 1,050$.

E.—A cross section of a small mature column. $\times 525$.






PLATE 53

Cronartium ribicola:



A drawing to show the intimate relation of the mycelium of the parasite to the host cells. *pp*, Phloem parenchyma; *sc*, sieve cells; *c*, cambium cells; *t*, tracheids; *rdp*, resin-duct parenchyma. The number of haustoria represented as entering the cells in this drawing is not abnormally large. The drawing was made from two serial sections from the same area by means of a projection apparatus and a camera lucida and has been diagrammatized only so far as was necessary to bring some of the elements to a proper level for drawing. In a few cases nuclei, which were not present in the sections, were supplied for both host and parasite cells. Note the different shapes and sizes of the haustoria and the general character of the hyphæ lying between the cells. $\times 500$.

PLATE 54

Cronartium ribicola:

A.—A drawing of an infected 12-year-old main stem. The infection entered the main stem along the small branch, the stub of which is shown at the right of the figure. *a*, The advancing edge of the infection; *b*, the pycnial area. The black dots are the pycnial spots; *c*, the æcial area on which the bark is cracked and broken. In another season the æcial area would spread over the pycnial area (*b*), and the pycnial area would be advanced as far as the boundary (*a*) under normal conditions. The boundary (*a*) would be proportionately advanced also. The specimen from which the drawing was made was collected in August, 1917. $\times \frac{1}{4}$.

B.—Drawing of a section through part of a young æcium showing the relation of the fertile cells with their denser protoplasmic contents to the overlying sterile cells, in which the cytoplasm and nuclei have begun to go to pieces. The manner in which the adjacent host cells are forced apart by the fungus cells is also shown. *dc*, decomposition products in the host cells; *hc*, host cell wall; *sc*, sterile cells; *fc*, fertile cells; *h*, haustorium. The elliptical bodies in the host cells represent starch grains. $\times 400$.






PLATE 55

Cronartium ribicola:

A.—A drawing of a median section through a very young uredinium. Note that the fungus cells are closely compacted against the underside of the host epidermal cells, and that the binucleate cells adjacent to the epidermal cells are oriented with their long axis more or less perpendicular to the epidermis. The section from which the drawing was made was cut just to one side of a stoma. See text for further explanation. $\times 500$.

B.—A drawing of a median section through a young uredinium, somewhat older than that illustrated in figure A. *a*, A peridial cell; *b*, a young urediniospore; *c*, a cell which is homologous to the stalk cell of an older sorus; *d*, a basal cell. See text for further explanation. $\times 500$.

C.—A drawing of a median section through a small mature uredinium. *a*, A binucleate basal cell; *b*, a tetranucleate basal cell just before the urediniospore initial is cut off; *c*, a secondary urediniospore initial which has just been cut off from the basal cell; *d*, the urediniospore initial has divided into a urediniospore and a stalk cell; *e*, a mature urediniospore which is still connected with the basal cell by a collapsed stalk cell; *f*, crushed epidermal cells of the host; *g*, the bank of parenchyma-like cells which encircle the sorus. The cells of this group, which lie next to the epidermis, are homologous to the thickened peridial cells which overlie the greater part of the uredinium. Note the comparatively small opening in the peridium. The drawing has been slightly diagrammatized for the sake of completeness by combining some of the features from two sections. Compare Plate 51, B. $\times 500$.




PLATE 56

Cronartium ribicola:

A.—A drawing of the cell relations near the edge of an æcium to illustrate the formation of the multilayered æcial peridium. *a*, A young æciospore; *i*, an intercalary cell; *b*, the basal cell of the chain; *φ*, the potential æciospores which are being modified into peridial cells. Note the wall markings and the long degenerating intercalary cells. $\times 400$.

B.—A drawing of a section through a mature peridium, taken from the same series as the photomicrograph in Plate 50, B. $\times 400$.

C.—A drawing of a short mature telial column in which the teliospores (*a*) have germinated, producing promycelia and sporidia (*b*). See Plate 57 for details of the process. $\times 170$. Drawn by Miss Minnie W. Taylor.






PLATE 57

Cronartium ribicola:

- A.—A mature teliospore from the tip of a column. $\times 850$.
- B.—A mature teliospore from the side of a column. $\times 850$.
- C.—A mature teliospore from the side of a column. $\times 850$.
- D.—Five germinating teliospores from a longitudinal section of a column. The nucleus in the promycelium from the upper teliospore is in the telophase of the primary division. $\times 850$.
- E.—A germinating teliospore. The nucleus is about to pass through the germ pore into the promycelium. $\times 850$.
- F.—Early prophase of the primary division in the promycelium. $\times 1,700$.
- G.—Late prophase of the primary division in the promycelium. The chromatin is in the form of complex tangle. $\times 1,700$.
- H.—Early anaphase of the primary division in the promycelium. Eleven chromosomes are visible. $\times 1,700$.
- I.—Later anaphase than that shown in figure H. Sixteen chromosomes visible. $\times 1,700$.
- J, K.—Two anaphase stages of the primary division. The chromosomes are well advanced toward the poles. $\times 1,700$.
- L.—Late anaphase of the primary division. The chromosomes are condensing into deeply staining clumps, but the individuals are still discernible in some cases. $\times 1,700$.
- M.—End of the anaphase of the primary division. The fibres connecting the two chromatin groups are drawn to a thin fading strand. Note the radiations in the cytoplasts in this and the preceding figure. $\times 1,700$.
- N.—The 2-celled promycelium. $\times 850$.
- O.—Metaphase of the second division. $\times 1,700$.
- P.—Telophase of the second division. Peculiar cytoplasmic radiations run from the reorganizing nuclei. $\times 1,700$.
- Q.—The reorganizing nuclei after the second division. $\times 1,700$.
- R.—The completed promycelium. Each nucleus shows a small nucleolus and a definite centrosome. $\times 850$.
- S.—Surface view of a germinating promycelium. $\times 850$.
- T.—The tip cell of a germinating promycelium. The nucleus exhibits definite polarization. $\times 1,700$.
- U.—A little later stage than the last. The nucleus is passing into the sterigma on its way to the sporidium. $\times 850$.
- V.—Surface view of a tip cell of a promycelium bearing a sterigma and a nearly mature sporidium. $\times 850$.
- W.—A mature sporidium. The papilla marks the point of attachment of the sterigma. $\times 850$.
- X, Y, Z.—Steps in the germination of the sporidia. $\times 850$.
- AA.—Sectional view of a mature sporidium. The nucleus shows polarization. $\times 1,700$.
- BB.—Sectional view of a germinating sporidium. The nucleus appears to be moving toward the germ tube and preparing to divide. $\times 1,700$.
- CC, DD.—Two stages in the formation of secondary sporidia. $\times 850$.
- EE.—Midanaphase of the division of the sporidium nucleus. $\times 1,700$.
- FF.—Late anaphase of the same. $\times 1,700$.
- GG.—Sectional view of a binucleate sporidium. $\times 1,700$.
- HH.—Two cells from the vegetative mycelium in the pine. $\times 1,700$.
- II.—A definitely polarized nucleus from the vegetative mycelium in the pine, located just beneath the fertile layer of the young aëcium. $\times 1,700$.

PLATE 58

Cronartium ribicola:

A.—The elements of the pycnium. The cells at the base are almost empty. Above them are the short branching trunks which bear the sporophores. *a*, A sporophore; *b*, pycniospores in sectional view; *c*, the cytoplasm is constricted just beneath the spore; *d*, the nucleus is dividing. $\times 1,700$.

B.—An active thin-walled haustorium from a pine host cell. $\times 1,700$.

C, D, E.—Old haustoria. C and D have basal cuplike sheaths and tip sheaths. E is completely inclosed in a thick sheath. $\times 850$.

F.—Telophase of division of one of the cells of the fertile layer to form a sterile cell. $\times 1,700$.

G.—A newly formed sterile cell. $\times 1,700$.

H.—A large polarized nucleus from the fertile layer. $\times 1,700$.

I.—An aëial basal cell resulting from the fusion of two adjacent cells of the fertile layer. Compare the size of the nuclei with that of the nucleus shown in Pl. 57, II. $\times 1,700$.

J.—A diagram of a basal cell resulting from the fusion of two cells from different levels. $\times 850$.

K.—A diagram of a trineucleate irregular basal cell from the tip of which a trineucleate aëiospore initial has been cut off. $\times 850$.

L.—A diagram of part of an irregular compound fusion cell. $\times 850$.

M.—A basal cell with the nuclei in early prophase. The centrosome in the left nucleus has apparently divided. *b*. An aëiospore initial cell with the nuclei in midanaphase. $\times 1,700$.

N.—Part of a basal cell. The nuclei in prophase. $\times 1,700$.

O.—A later stage than the preceding. The spindle and centrosomes are just visible in the upper nucleus. $\times 1,700$.

P.—Metaphase of the division in the basal cell. A centrosome is evident at each of the poles of the spindles. The chromatin is condensed into chromosomes. The nucleoli have moved off and begun to fade. $\times 1,700$.

Q.—Early anaphase, a little later than the stage in figure P. $\times 850$.

R.—Midanaphase of the division. The chromosomes are moving toward the poles along the outside of the spindle. $\times 1,700$.

S.—A later stage of anaphase than in figure R. $\times 1,700$.

T.—Final anaphase. The chromatin is condensing in two groups at each pole. $\times 1,700$.

U.—Telophase. The two groups for each pole are still distinct. $\times 1,700$.

V.—Telophase. The two groups at each pole have condensed to single masses. The wall which will separate the aëiospore initial from the basal is beginning to form at *a*. $\times 1,700$.


W.—Prophase of the division in the aëiospore initial. $\times 1,700$.

X.—Final telophase of the same. $\times 1,700$.

Y.—An aëiospore chain in section view. *a*, The basal cell; *b*, an aëiospore initial, *c*, an intercalary cell; *d*, a young aëiospore. The nuclei in the upper intercalary cells are degenerating. $\times 850$.

Z.—A large mature aëiospore in surface view. $\times 850$.

AA.—A nucleus from a mature aëiospore. $\times 1,700$.






PLATE 59

Cronartium ribicola:

- A.—Germinating aëciospores. $\times 425$.
- B.—Sectional view of the aëciospore wall showing the manner in which the germ tube is constricted. $\times 850$.
- C.—A binucleate cell from the mycelium in a leaf of *Ribes* sp. The nuclei are polarized as in the uninucleate mycelium. $\times 1,700$.
- D.—A binucleate haustorium from a host cell of *Ribes* sp. $\times 1,700$.
- E.—A uredinal basal cell. The nuclei are in prophase. $\times 1,700$.
- F.—Metaphase of the primary division in the basal cell. The chromosomes are visible as distinct units. $\times 1,700$.
- G.—Early anaphase of the same division. $\times 1,700$.
- H.—A later stage of the anaphase. $\times 1,700$.
- I.—Final anaphase. The two groups at each pole are not clearly distinct in this figure. $\times 1,700$.
- J.—A binucleate urediniospore initial. The nuclei are polarized. $\times 1,700$.
- K.—Metaphase or early anaphase of the division in the initial. $\times 1,700$.
- L.—Late anaphase in the initial. The chromatin is condensing in two groups at each pole. The second nucleus in the basal cell has been cut away. $\times 1,700$.
- M.—Late anaphase group from the initial, for comparison with figure L. $\times 1,700$.
- N.—A basal cell bearing a stalk cell surmounted by a nearly mature urediniospore (a), and a secondary urediniospore initial (b). The nuclei in the latter are in prophase. $\times 850$.
- O.—Metaphase of the secondary division in the basal cell, preparatory to the formation of a secondary urediniospore initial. $\times 1,700$.
- P.—A mature urediniospore. $\times 850$.
- Q.—A germinating urediniospore in sectional view. $\times 850$.
- R.—A germinating urediniospore. $\times 425$.
- S.—A telial basal cell. Both nuclei show centrosomes and polarization phenomena. $\times 1,700$.
- T.—Metaphase of the division in the telial basal cell. $\times 1,700$.
- U.—Late anaphase of the same. $\times 1,700$.
- V.—Final anaphase of the same. $\times 1,700$.
- W.—Telophase of the same. $\times 1,700$.
- X.—A diagram of a telial unit column. Two young binucleate teliospores surmount the basal cell. $\times 850$.
- Y.—The two nuclei of the young teliospore just previous to fusion. Note the centrosomes. $\times 1,700$.
- Z.—The two nuclei in the process of fusion. Two nucleoli and two centrosomes are still visible. $\times 1,700$.
- AA.—The large fusion nucleus. The chromatin is in heavy strands. $\times 1,700$.
- BB.—The fusion nucleus, slightly decreased in size. The chromatin strands are finer than in the preceding stage. $\times 1,700$.
- CC.—The mature fusion nucleus. The chromatin is in the form of granules. $\times 1,700$.

FURTHER DATA ON THE SUSCEPTIBILITY OF RUTACEOUS PLANTS TO CITRUS-CANKER¹

By H. ATHERTON LEE

Assistant Pathologist, Fruit-Disease Investigations, Bureau of Plant Industry, United States Department of Agriculture

INTRODUCTION

In the present paper results are given of field observations and inoculations with *Pseudomonas citri* upon plants belonging to genera more or less related to Citrus. These results show a wide range of hosts for Citrus-canker, and the possibility of lesions upon plants other than *Citrus* spp. serving as sources of new infection becomes emphasized in Citrus-canker eradication work.

In the present investigation the findings of Jehle² as to the susceptibility of *Chalcas (Murraya) exotica* Millsp., were corroborated and 23 other species representing 20 genera of the Rutaceae were studied.

It seems inadvisable to present the inoculation data here, since over 2,000 inoculations were made. Table I presents the data in a very much condensed form, and the illustrations show a few of the positive results. To anyone interested the complete inoculation data are available at the Bureau of Plant Industry, Washington, D. C. The inoculations and the controls were dried as herbarium specimens and will also be available for examination.

In making the inoculations an infusion of the organism was painted upon the leaf blade, midrib, petiole, or stem, as the case might be, with a small camel's-hair brush, and then the tissue was punctured through the coating of infusion with a needle. The inoculated twig was maintained in a moist condition by wrapping it in paraffin paper, including

¹ The investigations here outlined have been carried on largely at the Lanao Experiment Station of the Division of Plant Industry of the Philippine Bureau of Agriculture. Thanks are due to Mr. A. M. Burton, former Chief, and to Mr. S. Apostol, present Chief of this Division, as well as to Mr. F. G. Galang, Superintendent of the Lanao station. The Citrus collection of the College of Agriculture at Los Baños was also placed at the use of the writer through the courtesy of Dr. C. F. Baker, Dean of the College.

Many of the plants studied, belonging to genera closely related to Citrus are growing in the experimental plots of Mr. W. T. Swingle, Physiologist in Charge of Crop Physiology and Breeding Investigations, Bureau of Plant Industry, United States Department of Agriculture, for use in the breeding of canker-resistant Citrus fruits and canker-resistant stocks. Permission to use this material has greatly facilitated the work. Thanks are also due to Mr. Swingle for many helpful suggestions. Many plants related to Citrus grow wild in the Philippine Islands. Prof. E. D. Merrill, Botanist of the Bureau of Science, made helpful suggestions as to such plants and identified many of them. His help has been of the greatest value, and is hereby gratefully acknowledged.

It would have been difficult, if not impossible, to carry on this work without the extensive laboratory facilities of the Bureau of Science so freely made available through the courtesy of Dr. A. J. Cox, Director.

² JEHL, R. A. SUSCEPTIBILITY OF NON-CITRUS PLANTS TO BACTERIUM CITRI. In *Phytopathology*, v. 7, no. 5, p. 339-344. 3 figs. 1917.

with the twig also a small piece of moistened cotton. Control needle punctures with tap water were made for each host plant, and the moist condition was also maintained with paraffin paper and moist cotton. Unless otherwise noted, the positive results observed here occurred only at needle punctures.

TABLE I.—Summary of results of inoculations on plants of Rutaceae

No.	Genus and species.	Inoculation No. for reference.	Result.	Remarks.
SUBFAMILY CITRATAE. (Not of tribe Citreae.)				
1	<i>Claucaena laniatum</i>	1046-1050.....	Positive.....	Negative for leaves, but positive on petioles and stems; inoculations slow in maturing.
2	<i>Chalcas exotica</i>	876-880.....	Weakly positive.....	Negative for leaves but positive on petioles and stems. Susceptibility only evidenced by slight swelling not evidenced in controls.
TRIBE CITREAE. SUBTRIBE Feroninae.				
3	<i>Feronia limonia</i>	41-50.....	Positive.....	Leaves as well as stems show positive results. Reaction takes place slowly.
4	<i>Feroniella lucida</i>	1361-1370.....do.....	Results obtained only for stem. Positive results obtained very readily.
5	<i>Aegle marmelos</i>	811-816.....	Negative.....	Inoculation attempts made repeatedly on all parts of plants; all negative; believed to be immune.
6	<i>Chaetochloa glutinosa</i>	61-70; 831.....	Positive.....	Results obtained very readily in 5 days. Many naturally occurring infections (Pl. 62).
7	<i>Balsamocitrus gabonensis</i>	1341-1350.....	Negative.....	This species is believed to be immune.
8	<i>Subtribe Lavanginae.</i> <i>Hesperethusa crenulata</i>	1291-1310.....	Positive.....	Results obtained quickly and readily for both leaves and stems (Pl. 61, A).
9	<i>Triphasia trifolia</i>	1071-1808.....	Negative.....	Stems and leaves negative.
10	<i>Paramignya longipedunculata</i>	771-790.....	Positive.....	Tissue yellow, with only appearance around punctures, not raised; no such coloration around control; true both for leaves and stems (Pl. 61, B).
11	<i>Severinia burifolia</i>	1381-1390.....	Negative.....	Believed to be immune; inoculated repeatedly under same conditions which gave positive results on other genera.
SUBTRIBE CITRINAE.				
12	<i>Citropsis schweinfurthii</i>	1371-1380.....	Positive.....	Readily positive both leaves and stem.
13	<i>Atalantia citrioides</i>	1331-1335.....do.....	Positive for leaves, causing a watery dark discoloration of tissue but no excrescence. Weakly positive on stem (Pl. 61, A).
14	<i>Atalantia disticha</i>	936-950.....	Weakly positive.....	Stem inoculations slightly swollen, not the case with controls. Leaf inoculations slightly discolored, not the case with controls. Very resistant at least.
15	<i>Eremocitrus glauca</i>	1416-1420.....do.....	Stem inoculations slowly positive.
16	<i>Fortunella hindsii</i>	1421-1480.....	Positive.....	Quickly positive on both leaves and stem (Pl. 61, B).
17	<i>Fortunella japonica</i>	1031-1040.....	Weakly positive.....	Leaf blade inoculations definitely negative; inoculations in midrib weakly positive and inoculations in stems slowly but clearly positive. Believed to be highly resistant.
18	<i>Microcitrus australis</i>	486-505.....	Positive.....	Quickly positive; cankers definite but much smaller than on <i>Citrus</i> spp.
19	<i>Microcitrus australatica</i>	1126-1130.....do.....	Stem inoculations show cankers; much smaller than on <i>Citrus</i> spp., however.

TABLE I.—Summary of results of inoculations on plants of Rutaceae—Continued

No.	Genus and species.	Inoculation No. for reference.	Result	Remarks.
RUTACEAE OTHER THAN CITRATAE.				
30	<i>Toddalia asiatica</i>	1401-1410.....	Positive.....	Leaves show no excrescences, but a yellowing of tissue not found in control punctures. Stem inoculations are swollen and the tissue is blackened.
21	<i>Xanthoxylum rhetsa</i>	631-690.....	Negative.....	Leaves, petioles, and stems negative.
22	<i>Evodia ridleyi</i>	2113-2127.....	Positive.....	Both leaves and stem produce light brown eruptions very definite and similar to those on <i>Citrus</i> spp. (Pl. 65). There is also evidence that stomatal infections took place.
23	<i>Evodia latifolia</i>	291-300.....	do.....	Leaves clearly negative but stem tissues show excrescences of a brown color similar to those produced on <i>Citrus</i> spp.; lesions very large.
24	<i>Melicope triphylla</i>	1526-1540.....	do.....	Leaves clearly negative; stem tissue shows excrescences of brown color similar to those produced on <i>Citrus</i> spp.

Field observations corroborating these inoculation data have been made in many cases. Thus, in South China *Severinia buxifolia* occurs naturally, exposed in a number of instances to infection from Citrus-canker in near-by orchards. In no instances were lesions in any way resembling those of Citrus-canker found. *Aegle marmelos* and *Triphasia trifolia* are cultivated at Lamao in the Philippine Islands, surrounded by nursery rows of Citrus trees heavily infected with canker. In no case have lesions similar to those of canker been found on these hosts.

Chalcas exotica is a commonly-grown ornamental in Manila; nothing at all resembling Citrus-canker has ever been found occurring naturally upon it, although sources of infection are in some cases closely present.

Xanthoxylum rhetsa and *Alalantia disticha* occur naturally at Lamao in places where infection with Citrus-canker would be easily possible. No lesions similar to canker have been found on these plants.

Fortunella japonica occurs in orchards at Lamao and also at Los Banos, Philippine Islands. No naturally-occurring cankers have ever been seen on this host. Wolf¹ reports Citrus-canker on kumquats, but does not mention what species were under observation. Swingle² also reports canker on kumquats in Japan.

Fortunella japonica, although susceptible to Citrus-canker under the most optimum conditions, should nevertheless be regarded as highly resistant, closely approaching immunity.

On the other hand, *Fortunella hindsii* occurring naturally in South China has been observed frequently with heavy canker infection. Plants

¹ Wolf, Frederick A. CITRUS-CANKER. *In* Jour. Agr. Research, v. 6, no. 2, p. 70. 1916.

² U. S. DEPARTMENT OF AGRICULTURE. CITRUS-CANKER IN PHILIPPINES. *In* U. S. Dept. Agr. Dept. Circ., v. 1, no. 1, p. 8. 1915.

of this species were found by Prof. G. W. Groff and the writer near the summits of mountains in Kwangtung Province at an altitude of about 1,500 feet. These mountains are heavily eroded and peculiar in having their sides barren and almost entirely free from growth. Plants of *F. hindsii* at the tops of these mountains were therefore very much isolated from commercial Citrus plantings and other sources of canker infection; nevertheless, in almost all cases the species was heavily infected with Citrus-canker. The writer later found plants of the same species upon Victoria Peak, Hongkong, a mountain of very similar type. It is possible that further facts may show that this plant is an original wild host from which Citrus-canker has spread to cultivated species.

Chaetospermum glutinosa occurs both naturally and cultivated at Lamao, and naturally occurring cankers are abundant upon such plants. The susceptibility of *C. glutinosa* to canker is easily greater than that of the sweet orange (*Citrus sinensis*) in the Philippines.

Glycosmis pentaphylla occurs naturally in Kwangtung Province, China, in places where infection from Citrus-canker would be easily possible. No naturally occurring cankers were observed.

Of the positive results obtained in the foregoing tabulated species, *Pseudomonas citri* has been reisolated from *Clauцена lانسium*, *Feronia limonia*, *Feroniella lucida*, *Chaetospermum glutinosa*, *Hesperethusa crenulata*, *Paramignya longipedunculata*, *Citropsis schweinfurthii*, *Atalantia citrioides*, *Fortunella hindsii*, *Microcitrus australasica*, *M. australis*, *Toddalia asiatica*, and *Evodia ridleyi*. Such isolations have been reinoculated on foliage of *Citrus grandis*, and have given positive results in each case. In those positive results not listed as having the organism reisolated the material was collected and dried in the field, where laboratory facilities were not available.

The statements made above as to immunity, since they are based on lack of infection after inoculation with a dense infusion of the causal organism under the most favorable conditions for infection, are probably more substantial than claims made for absence of infection under natural conditions.

The most noteworthy feature of the inoculations is the susceptibility of such very distant relatives as *Evodia ridleyi*, *E. latifolia*, and *Melicope triphylla*. In these cases the cankers are by no means weakly produced, but form quickly, with a decided swelling of the tissue, which later erupts as on species of Citrus. On *E. ridleyi* the results were evident in two weeks. *Pseudomonas citri*, therefore, is not closely limited to Citrus spp., but has a very wide range of host plants within the family Rutaceae.

Severinia buxifolia, *Aegle marmelos*, and *Balsamocitrus gabonensis* produce no reaction whatever when inoculated with *Pseudomonas citri*. It is believed that these species may be safely called immune to Citrus-canker; this is especially noteworthy, since they are all close relatives of the genus Citrus.

The evidence is that *Xanthoxylum rhetsa* and *Triphasia trifolia* are immune, while *Chalcas* (*Murraya*) *exotica*, *Atalantia disticha*, and *Fortunella* (*Citrus*) *japonica* show positive results only under the most favorable circumstances for infection.

It is noteworthy that some of the immune and highly resistant species possess thick, coriaceous brittle leaves—for example, *Severinia buxifolia*, *Atalantia disticha*, *Fortunella japonica*, and *Chalcas exotica*. This suggests the possibility that resistance to Citrus-canker may be influenced in some measure by histological or morphological differences.

SUMMARY

(1) Inoculation tests with *Pseudomonas citri* upon 24 species representing 20 genera of the family Rutaceae show that 19 of the species are susceptible in greater or less degree. It thus appears that Citrus-canker is not closely limited to the genus *Citrus*, but has a wide range of hosts among the Rutaceae.

(2) *Severinia buxifolia*, *Aegle marmelos*, and *Balsamocitrus gabonensis*, all close relatives of *Citrus*, may safely be called immune to Citrus-canker. *Xanthoxylum rhetsa* and *Triphasia trifolia* seem to be immune.

(3) *Chalcas* (*Murraya*) *exotica*, *Atalantia disticha*, and *Fortunella* (*Citrus*) *japonica*, also closely related to the genus *Citrus*, are strongly resistant to Citrus-canker.

(4) *Clauцена lانسium*, *Feronia limonia*, *Feroniella lucida*, *Chaetoppermum glutinosa*, *Hesperethusa crenulata*, *Paramignya longipedunculata*, *Citropsis schweinfurthii*, *Atlantia citrioides*, *Fremocitrus glauca*, *Fortunella hindsii*, *Microcitrus australis*, *M. australasica*, *Toddalia asiatica*, *Evodia ridleyi*, *E. latifolia*, and *Melicope triphylla*, of different relationships to the genus *Citrus*, all produce positive results when inoculated with *Pseudomonas citri*, at needle punctures. Of these *Clauцена lانسium* and *Feronia limonia* develop infection very slowly, the others fairly quickly.

(5) *Chaetoppermum glutinosa* shows naturally occurring infections of Citrus-canker and in the Philippines its susceptibility is easily greater than that of the sweet orange (*Citrus sinensis*). *Fortunella hindsii* occurs naturally in South China, very much isolated from sources of Citrus-canker infection. The abundance of cankers found on such trees gives rise to the theory that this species may have been an original wild host from which Citrus-canker spread to cultivated species.

PLATE 6o

Naturally occurring Citrus-canker lesions on leaves of *Chaetospermum glutinosa*.
Natural size.

(666)







PLATE 61

A.—*Hesperethusa crenulata*, showing discolorations resulting from inoculations with *Pseudomonas citri*. Natural size.


B.—*Paramignya longipedunculata*, showing discolorations around punctures made with *P. citri* on leaves. Such discolorations did not develop at punctures made with river water. Natural size.

PLATE 62

A.—*Atalanta citrioides*, showing positive results following inoculation with *Pseudomonas citri*. Natural size.

B.—*Fortunella hindsii*, showing results of inoculation with rain water on leaves. Natural size.

C.—*Fortunella hindsii*, showing results of inoculation with *P. citri* on leaves. Natural size.






PLATE 63

Evodia ridleyei:

A.—Stem inoculated with tap water.

B, C.—Two twigs inoculated with *Pseudomonas citri*. Natural size.

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
25 CENTS PER COPY
△

INDEX

	Page		Page
Acetate, calcium, method used in determining soil acidity.....	322-329	<i>Agropyrum</i> —Continued	
<i>Achorion Schönlensis</i> , causal organism of favus.....	415-418	<i>repens</i> —	
Acid—		source of <i>Puccinia graminis</i> in bridging-host experiment.....	328
amino, in normal and blighted spinach.....	385-404	susceptibility to <i>Puccinia graminis tritici</i>	323
hydrochloric, used in—		<i>tritici</i> —	
analyzing greensand.....	485	susceptibility to <i>Puccinia graminis tritici</i>	324
determining water-soluble nitrogen in spinach.....	397-404	used in bridging-host experiment.....	331-340
hydrofluoric, used in analyzing greensand.....	489	<i>lenerum</i> —	
lactic, in alfalfa silage.....	581	source of <i>Puccinia graminis tritici</i> transferred to semicongenial hosts.....	349
nitric, used in experiments with plowsole.....	508	susceptibility to <i>Puccinia graminis tritici</i>	324
nitrohydrochloric, used in experiments with plowsole.....	507	<i>Agrostis</i> —	
phosphomolybdic, used in determining nitrogen in spinach.....	399-404	<i>alba</i> , source of <i>Puccinia graminis agrostis</i> transferred to uncongenial hosts.....	345
phosphotungstic, used in—		<i>stolonifera</i> —	
determining basic nitrogen in spinach.....	399-404	source of <i>Puccinia graminis</i> in bridging-host experiment.....	328
titration of silage extract.....	130-131	source of <i>Puccinia graminis agrostis</i> transferred to uncongenial hosts.....	345
sulphuric, used in—		Albuminoid nitrogen in silage.....	129-131
damping-off experiment.....	526	Alcohol—	
determining basic nitrogen in spinach.....	399-404	in silage—	
Acidity of—		caused by zymase.....	590
alcoholic extract of silage.....	116-125	released by enzymic and yeast action.....	589-590
dioxigen.....	140-141	tolerance of larvae of <i>Trichinella spiralis</i> to.....	477-479
hydrogen peroxid.....	140-141	to.....	477-479
plant sap, changes during freezing.....	58-101	Alcoholic extract of silage—	
seeds, effect on catalase activity.....	141-143	acidity.....	116-125
silage.....	574, 581	method of making.....	115
soil, effect of moisture.....	321-329	Alfalfa—	
water extract of silage.....	116-125	effect on plowsole.....	507
Acid-producers in alfalfa silage.....	574-592	hay—	
Acids—		digestibility.....	371, 383
diamino, in healthy and blighted spinach.....	399-404	methane fermentation.....	374-375
tolerance of larvae of <i>Trichinella spiralis</i> to.....	477-479	net energy value.....	269-286
<i>Acromoniella</i> sp., fungus isolated from <i>Ipomoea batatas</i>	364	silage—	
<i>Actinomyces</i> sp., fungus isolated from <i>Ipomoea batatas</i>	364	bacteria in.....	571-592
Adams, J. M. R., et al. (paper): Sweet-Potato Storage-Rots.....	317-368	comparison with sweet-clover silage.....	113-113
<i>Adelphocoris rapidos</i> , attacking clover.....	181	Alkali—	
Adobe soil, inorganic colloids.....	516	effect of soil factors on toxicity.....	387-319
<i>Aephe marmelos</i> , immunity to <i>Pseudomonas citri</i>	662-665	salts, relative toxicity.....	315-318
After-ripening of seeds, effect on—		Alleys, effect on border rows in plot tests.....	257-261
catalase activity.....	138, 149-159	<i>Alisma</i> spp., nature of dormancy of seeds.....	138
oxidase activity.....	163-166	<i>Alternaria</i> sp., isolated from—	
<i>Agrotis mancur</i> , larvae, effect of hydrocyanic-acid gas on.....	134-135	dead conifers.....	548
<i>Agropyrum</i> —		<i>Ipomoea batatas</i>	348, 355
<i>convivum</i> , source of <i>Puccinia graminis tritici</i> transferred to semicongenial hosts.....	342	Aluminum—	
<i>crinitum</i> —		in mulch, plowsole, and subsoil.....	508-509
source of <i>Puccinia graminis</i> in bridging-host experiment.....	332	oxid in—	
source of <i>Puccinia graminis tritici</i> transferred to semicongenial hosts.....	342	mulch, plowsole, and subsoil.....	508-526
		normal and blighted spinach.....	371-375
		soil, effect of drying.....	325-358
		<i>Amaranthus</i> —	
		<i>blaboides</i> , catalase content of seeds.....	144, 153-154
		<i>gracilis</i> , catalase content of seeds.....	153-154
		<i>retrofractus</i> , catalase content of seeds.....	144, 149, 153-154
		spp. nature of dormancy of seeds.....	138

	Page		Page
<i>Ambrosia</i> —		<i>Aspergillus</i> —	
<i>artemisiacifolia</i> , catalase and oxidase content		<i>niger</i> , fungus isolated from <i>Ipomoea batatas</i>	364
of seeds.....	138	sp., not cause of damping-off in conifers.....	549
<i>trifida</i> , catalase and oxidase content of seeds	138	<i>Atlantia</i> —	
Amids, acid, in normal and blighted spinach.....	385-404	<i>citrioides</i> , susceptibility to <i>Pseudomonas citri</i>	662-665
Amino—		<i>didicha</i> , susceptibility to <i>Pseudomonas citri</i>	662-665
acids in normal and blighted spinach.....	385-404	<i>Aucuba japonica</i> , production of injected areas due to freezing.....	86
nitrogen in—		Austrian pine. See <i>Pinus austriaca</i> .	
alfalfa silage.....	581-590	<i>Avena sativa</i> —	
hardened and nonhardened cabbage.....	97	reported bridging host for transfer of timothy rust.....	222
silage extract.....	125-132	used in bridging-host experiments.....	226-249
Ammonia—		Azotobacter, effect of carbon disulphid and toluol on.....	601-614
effect of carbon disulphid and toluol on accumulation in soil.....	603-614	Bacteria, nitrogen-fixing, effect of carbon disulphid and toluol.....	601-614
in alfalfa silage.....	586-590	Bacteriological Studies on Alfalfa Silage (paper).....	571-592
in hardened and nonhardened cabbage.....	97	<i>Bacterium lactrymans</i> , causal organism of angular-leaftspot of cucumber.....	201-220
Ammonia nitrogen in healthy and blighted spinach.....	398-404	<i>Bactroera cucurbitae</i> —	
Ammonium—		host of <i>Opius fletcheri</i>	456-457
carbonate, used in experiments with plowsole.....	505	host of <i>Tetrastichus giffardianus</i>	454-458
hydrate, used in experiments with plowsole.....	505	Balsam-apple. See <i>Momordica balsamina</i> .	
<i>Amygdalus persica</i> , nature of dormancy of seeds.....	138	<i>Balsamocitrus gabonensis</i> , immunity to <i>Pseudomonas citri</i>	561-665
<i>Andropogon halepensis sudanensis</i> , catalase and oxidase content of seeds.....	137-174	Balsam-pear. See <i>Momordica charantia</i> .	
Angular-Leaftspot of Cucumber: Dissemination, Overwintering, and Control (paper).....	201-220	Barberry. See <i>Berberis</i> spp.	
Angular-leaftspot of cucumber—		Barium—	
dissemination by—		chlorid, used in determining nitrogen in spinach.....	400-404
drainage water.....	211	hydroxid, used in determining nitrogen in spinach.....	399-404
insects.....	212-213	phosphotungstate, used in determining nitrogen in spinach.....	399-404
pickers.....	211-212	Harley. See <i>Hordeum</i> spp.	
rain and wind.....	208-210	Basic nitrogen in healthy and blighted spinach.....	399-404
effect of—		Basswood. See <i>Tilia americana</i> .	
desiccation.....	203-204	Beach, B. A. and Halpin, J. C. (paper): Observations on an Outbreak of Favus.....	415-418
germicides.....	205	Beef cattle, crosses with dairy breeds.....	1-57
fruit infection.....	208	Beetle, June green. See <i>Colinus nitida</i> .	
host plants.....	205-206	Begonia, production of injected areas due to freezing.....	86
stomatal infection.....	206-208	<i>Berberis</i> spp., reported bridging host for <i>Puccinia bovaliformis</i>	222
thermal death point.....	204-205	Bermuda grass. See <i>Capriola dactylon</i> .	
overwintering in—		Bestill. See <i>Thevetia nerifolia</i> .	
insects.....	214	<i>Beta vulgaris</i> , effect of farm manure on.....	494-503
seeds.....	214	Biologic forms of <i>Puccinia graminis</i> , plasticity of.....	221-249
soil.....	213	Biology of fruit-fly parasites in Hawaii.....	419-466
remedial measures—		Black, Otis F., et al. (paper): Physiological Studies of Normal and Blighted Spinach.....	369-408
resistant varieties.....	217	Blackrot—	
sanitation.....	217	Java, of <i>Ipomoea batatas</i>	347-349
seed treatment.....	218	of <i>Ipomoea batatas</i> —	
spraying.....	217-218	description.....	344-346
Ant. See <i>Phicidole megacephala</i> .		susceptible varieties.....	346
Armstrong, Henry Prentiss, and Fries, J. August (paper): Net Energy Values of Alfalfa Hay and of Starch.....	269-286	Blacktop, type of damping-off.....	545-558
Armstrong, A. C., and Hayes, H. K. (paper): Experiments in Field Technic in Plot Tests.....	251-262	Black-walnut worm. See <i>Dalania integririma</i> .	
Artschwager, Ernst F. (paper): Histological Studies on Potato Leafroll.....	559-570	Bluegrass. See <i>Poa pratensis</i> .	
Ash—			
digestibility in alfalfa hay and starch.....	271-282		
in normal and blighted spinach.....	371-375		
in dry matter of feeding stuffs.....	270		
Asparagin, used in determining water-soluble nitrogen in spinach.....	397-404		

	Page		Page
Bordeaux mixture, used against <i>Bacterium</i>		Carbohydrate changes during hardening	
<i>lachrymans</i>	217-218	against frost injury.....	95-97
<i>Botryorhiza hippocrateae</i> , difference from <i>Cro-</i>		Carbohydrate production in healthy and	
<i>narium rubicola</i>	627-628	blighted spinach.....	381-384
<i>Botrytis</i> —		Carbohydrates, effect on alfalfa silage.....	584-590
<i>cinerea</i> —		Carbon—	
cause of damping-off in conifers.....	547-556	digestibility in alfalfa hay and starch.....	271-284
fungus isolated from <i>Ipomoea batatas</i>	357-357	dioxide in—	
<i>douglasii</i> . Syn. <i>Botrytis cinerea</i> .		gaseous excretion.....	273, 284
<i>vulgaris</i> . Syn. <i>Botrytis cinerea</i> .		silage, released by enzymic and yeast	
Bouyoucos, George J., and McOwl, M. M.		action.....	589-590
(paper): Determining the Absolute Salt		disulphid, effect on nitrogen-fixing organ-	
Content of Soils by Means of the Freezing-		isms in soil.....	601-614
Point Method.....	331-336	in dry matter of feeding stuffs.....	270
<i>Brassica</i> —		in gaseous excretion.....	273, 284
<i>napus</i> , injected areas due to freezing.....	87	Carbonate—	
<i>oleracea capitata</i> , frost injury in.....	86	ammonium, used in experiments with	
Brier roses, resistant to brown canker.....	591-600	plowsole.....	508
Brisket disease—		calcium—	
condition of heart.....	409-411	effect on plowsole.....	507
effect of—		in olive earth.....	484
altitude.....	411-414	sodium, effect on plowsole.....	507
feed.....	411-413	Carsner, Eubanks (paper): Angular-Leadspot	
Bromelin, effect on oxidase activity of seeds.....	169	of Cucumber: Dissemination, Overwinter-	
<i>Bromus tectorum</i> —		ing, and Control.....	201-200
susceptibility to <i>Puccinia graminis</i>	222,	Caryopsis of seeds, catalase activity of.....	146-147
225-229, 240-241		Cinein (ligesterin) in alfalfa silage.....	586-590
used in bridging-host experiment.....	228-249	Catalase and Oxidase Content of Seeds in	
Brown Canker of Roses, Caused by <i>Diaporthe</i>		Relation to Their Dormacy, Age, Vitality,	
<i>umbriana</i> (paper).....	593-600	and Respiration (paper).....	137-174
Brown rust of bromes. See <i>Puccinia dispersa</i> .		Catalase of seeds—	
Brown-tail moth. See <i>Euproctis chrysorrhoea</i> .		method of determining.....	139-144
<i>Bryanopsis laciniata</i> , host plant of <i>Bacterium</i>		solubility.....	160-164
<i>lachrymans</i>	206	Cattle, crosses between dairy and beef	
Bug, gray-damself. See <i>Reduvius ferox</i> .		breeds.....	1-57
Eggnell, H. H., et al. (paper): Physiological		Cattleya orchids, fumigation with hydro-	
Studies of Normal and Blighted Spinach.....	369-408	cyanic-acid gas.....	263-268
Bulgarian bacteria in alfalfa silage.....	581-590	Cells, diameter in various parts of seeds of	
Hyrophylum, production of injected areas		Johnson grass and crimson clover.....	145
due to freezing.....	86-88	<i>Cephalothecium</i> sp., fungus isolated from	
Cabbage. See <i>Brassica oleracea capitata</i> .		<i>Ipomoea batatas</i>	364
Calabash gourd. See <i>Lagenaria vulgaris</i> .		<i>Ceratitix capitata</i> in Hawaii, parasitism.....	419-466
Calcium—		<i>Ceratomyx fimbriata</i> . See <i>Sphaerocoma fim-</i>	
carbonate—		<i>briatum</i> .	
effect on plowsole.....	507	<i>Ceratostoma</i> sp., fungus isolated from <i>Ipomoea</i>	
in olive earth.....	484	<i>batatas</i>	364
used to neutralize hydrogen peroxid.....	142-143	<i>Clostridium</i> sp., not cause of damping-off in	
in mulch, plowsole, and subsoil.....	502-516	conifers.....	540-558
calciumphosphate crystals, formation in frozen		<i>Chaetosporium glutinosum</i> , susceptibility to	
spots of cabbage leaf.....	86-87	<i>Pseudomonas citri</i>	602-605
nitrate in solution used in experiments with		<i>Chaetoxanthia</i> , susceptibility to <i>Pseudomonas</i>	
greensand.....	486-492	<i>citri</i>	601-606
oxid in—		Charcoal-rot of <i>Ipomoea batatas</i>	351-354
greensand.....	484	Chemical changes in plants during hardening	
mulch, plowsole, and subsoil.....	513-516	against frost injury.....	95-97
normal and blighted spinach.....	371-375	Chemistry of Sweet-Clover Silage in Compari-	
soil, effect of drying.....	755-758	son with Alfalfa Silage (paper).....	143-154
sulphate, effect on plowsole.....	527	Cherry, French. See <i>Eugenia uniflora</i> .	
Calcium-acetate method, used in determining		Chlorid—	
soil acidity.....	324-329	barium, used in determining nitrogen in	
Cane sugar, effect on alfalfa silage.....	581-590	spinach.....	400-404
Canker, brown, of roses.....	593-600	mercuric—	
<i>Capnodium meridionale</i> , similarity to <i>Coleo-</i>		effect on <i>Bacterium lachrymans</i>	205
<i>sporium senecioides</i>	648	effect on oxidase activity of seeds.....	168-169
<i>Capriola dactylon</i> , catalase and oxidase con-		used in determining nitrogen in spin-	
tent of seeds.....	137-174	ach.....	397-404

	Page		Page
Chlorid—Continued.		Complement-fixation test, serum distributed	
potassium, used in experiments with green-		by multiple-pipette holder.....	615-618
sand.....	488	Condition of Fertilizer Potash Residues in	
silver, used in determining nitrogen in		Hagerstown Silty Loam Soil (paper)....	59-81
spinach.....	400-404	Conifers, diseases of seedlings.....	521-558
sodium—		Conifer seedlings, injury from—	
effect on plowsole.....	597	damping-off.....	521-558
used in alkali experiment.....	269-319	drouth.....	554-555
used in experiments with intestinal tri-		mechanical means.....	551-554
chine.....	468-482	whitespot.....	551
Chloroform, effect on alfalfa silage.....	590	Conner, S. D. (paper): Soil Acidity as Affected	
<i>Chrysophyllum monopyrenum</i> , host of puparia		by Moisture Conditions of the Soil.....	321-329
parasitized by <i>Diachasma fullawayi</i>	418	Contribution to the Biology of Fruit-Fly	
Cimer—		Parasites in Hawaii (paper).....	413-466
<i>antennirectus</i> . See <i>Miris dolabratus</i> .		Copper—	
<i>deses</i> . See <i>Miris dolabratus</i> .		sulphate, effect on <i>Bacterium lachrymans</i>	295
<i>dolabratus</i> . See <i>Miris dolabratus</i> .		sulphid, used in determining nitrogen in	
<i>frumentarius</i> . See <i>Miris dolabratus</i> .		spinach.....	400-404
<i>laevigatus</i> . See <i>Miris dolabratus</i> .		<i>Coriscus ferus</i> , natural enemy of <i>Leptoterma</i>	
<i>lateralis</i> . See <i>Miris dolabratus</i> .		<i>dolabrata</i>	194
<i>porrectus</i> . See <i>Miris dolabratus</i> .		Corn chop, effect on alfalfa silage.....	574-590
<i>recticornis</i> . See <i>Miris dolabratus</i> .		Corrodentia, found on imported orchids.....	293
<i>riparius</i> . See <i>Miris dolabratus</i> .		Corsecan pine. See <i>Pinus nigra potrettiana</i> .	
<i>V-flavum</i> . See <i>Miris dolabratus</i> .		<i>Corticium vagum</i> , cause of damping-off in con-	
Citrate, susceptibility to <i>Pseudomonas citri</i>	662-665	ifers.....	521-558
Citreae, susceptibility to <i>Pseudomonas citri</i>	662-665	<i>Cobitis nitida</i> , larvae, effect of hydrocyanic-	
Citrinae, susceptibility to <i>Pseudomonas citri</i>	662-665	acid gas on.....	134-136
Citron. See <i>Citrullus vulgaris</i> .		<i>Crotalaria</i> spp., nature of dormancy of seeds.....	138
<i>Citrullus vulgaris</i> —		Crimson clover. See <i>Trifolium incarnatum</i> .	
free from attacks of <i>Bacterium lachrymans</i> ...	206	Crocker, William, and Harrington, George T.	
host of organism virulent to <i>Ipomoea batatas</i>	347	(paper): Catalase and Oxidase Content of	
Citrus canker. See <i>Pseudomonas citri</i> .		Seeds in Relation to Their Dormancy, Age,	
Citrus—		Vitality, and Respiration.....	137-174
<i>grandis</i> , susceptibility to <i>Pseudomonas</i>		<i>Cronartium</i> —	
<i>citri</i>	664-665	<i>asclerobium</i> , resemblance to <i>Cronartium</i>	
<i>sinensis</i> , susceptibility to <i>Pseudomonas</i>		<i>ribicola</i>	657
<i>citri</i>	665	<i>complanata</i> , nature of injury on <i>Pinus syl-</i>	
<i>Citropsis schweinfurthii</i> , susceptibility to		<i>vestita</i>	649
<i>Pseudomonas citri</i>	662-665	<i>flaccidum</i> , parasite on <i>Paeonia officinalis</i> ...	648-649
Clays, potash in.....	707-713	<i>occidentale</i> , <i>zeila</i>	652
<i>Claviceps lanarium</i> , susceptibility to <i>Pseudom-</i>		<i>ribicola</i> —	
<i>monas citri</i>	662-665	c, tology.....	630-654
Clover—		infection of pine.....	624-625
crimson. See <i>Trifolium incarnatum</i> .		life history.....	620-624
red. See <i>Trifolium pratense</i> .		morphology—	
<i>Coffea arabica</i> , host of puparia parasitized by		<i>zeila</i>	630-634
<i>Diachasma fullawayi</i>	418	mycelium.....	626-629
Coffee. See <i>Coffea arabica</i> .		mycelia.....	629-630
Coleoptera, found on imported orchids.....	263	telia.....	637-639
Coleosporium—		uredinia.....	635-637
<i>senecionis</i> , similarity to <i>Capnodium menis-</i>		Crook-neck squash, Japanese. See <i>Cucurbita</i>	
<i>male</i>	645	<i>moschata</i> .	
<i>solaniginis</i> . Syn. <i>Coleosporium senecionis</i> var.		Crosses between dairy and beef cattle—	
<i>senecionis</i>	645	breeds in parent herd.....	
<i>senecionis</i> , fusion nucleus in teliospore.....	646	inheritance of—	
Coleus, production of injected areas due to		beet qualities.....	57-58
freezing.....	86	color.....	57-58
Collembola, found on imported orchids.....	293	milk qualities.....	57-58
Colley, Reginald H. (paper): Parasitism, Mor-		muzzle color.....	45-48
phology, and Cytology of <i>Cronartium ribi-</i>		polluted character.....	45-48
<i>cola</i>	613-650	switch color.....	47-48
Colloids, soil, relation to plowsoles.....	305-319	tongue pigment.....	24-27
<i>Colocasia esculenta</i> , host of rot-producing		white margin.....	24-27
organism virulent to <i>Ipomoea batatas</i>	347	Crude fiber—	
Colon organisms in alfalfa silage.....	574-579	digestibility in alfalfa hay and starch....	271-272
Colorimetric titration of silage extract.....	127-129	in dry matter of feeding stuffs.....	270

	Page	<i>Diachasma</i> —Continued.	Page
Cucumber—		<i>trypsi</i> —	
squirting. See <i>Momordica elaterium</i> .		life history.....	450-459
wild. See <i>Echinocystis lobata</i> .		natural enemy of Mediterranean fruit	
See <i>Cucumis sativus</i> .		fly.....	459-460
<i>Cucumis</i> —		Diamino acids in healthy and blighted spi-	
<i>acutangulus</i> , host plant of <i>Bacterium lach-</i>		nach.....	392-424
<i>rymans</i>	256	<i>Diaporthe</i> —	
<i>anguria</i> , host plant of <i>Bacterium lach-</i>		<i>batatas</i> , causal organism of dryrot of <i>Ip-</i>	
<i>rymans</i>	256	<i>moea batatas</i>	349
<i>dipacrus</i> , host plant of <i>Bacterium lach-</i>		<i>umbrina</i> , n. sp., cause of brown canker of	
<i>rymans</i>	256	roses.....	593-600
<i>grossulariiformis</i> , free from attacks of <i>Bac-</i>		control.....	598-599
<i>terium lachrymans</i>	256	cultural characters.....	598
<i>melo</i> —		description.....	598-599
free from attacks of <i>Bacterium lachrymans</i>		life history.....	598
var. <i>dudaim</i> , free from attacks of <i>Bac-</i>		Dicotyledons, parasitized by same damping-	
<i>terium lachrymans</i>	256	off parasite as conifers.....	515-518
var. <i>flexuosus</i> , free from attacks of <i>Bac-</i>		Dietz, H. F., and Sasser, E. R. (paper):	
<i>terium lachrymans</i>	256	Fumigation of Cattleya Orchids with Hy-	
<i>sativus</i> , host plant of <i>Bacterium lachrymans</i> .	256	drocyanic-Acid Gas.....	261-268
<i>Cucurbita</i> —		Dimorphism of <i>Miris dolabratus</i>	183-184
<i>maxima</i> , free from attacks of <i>Bacterium</i>		Dioxid—	
<i>lachrymans</i>	256	carbon, in silage, released by enzymic and	
<i>marchola</i> , free from attacks of <i>Bacterium</i>		yeast action.....	589-590
<i>lachrymans</i>	256	silica in—	
<i>pepo</i> —		mulch, ploysele, and subsoil.....	528-536
free from attacks of <i>Bacterium lachrymans</i>		normal and blighted spinach.....	371-375
var. <i>condensa</i> , free from attacks of <i>Bac-</i>		Dioxigen, used in determining catalase con-	
<i>terium lachrymans</i>	256	tent of seeds.....	140-144
var. <i>onifera</i> , free from attacks of <i>Bacterium</i>		<i>Diplodia</i> —	
<i>lachrymans</i>	256	<i>gossypina</i> , causal organism of rot in <i>Ip-</i>	
Cuprous oxid in intumescences on frozen		<i>moea batatas</i>	347
leaves.....	88	<i>maculosa</i> , causal organism of rot in <i>Ipomoea</i>	
Cyanid—		<i>batatas</i>	347
potassium, tolerance of larvae of <i>Trichinella</i>		<i>natalensis</i> , causal organism of rot in <i>Ipomoea</i>	
<i>spiralis</i> to.....	477-479	<i>batatas</i>	347
sodium—		<i>tubercula</i> , causal organism of Java blackrot	
effect on cattleya orchids.....	267-268	of <i>Ipomoea batatas</i>	347
used to generate hydrocyanic-acid gas.....	334	<i>zeae</i> , nonvirulent to <i>Ipomoea batatas</i>	348
<i>Dactylis glomerata</i> —		Disaccharids in hardened and nonhardened	
food plant of <i>Miris dolabratus</i>	183	cabbage.....	95-96
reported bridging host for transfer of		<i>Discula pini</i> , not cause of damping-off in	
timothy rust.....	322	conifers.....	550
source of <i>Puccinia graminis philipponensis</i>		Diseases of conifer seedlings.....	521-528
transferred to senicongential hosts.....	244	Disulphid, carbon, effect on nitrogen-fixing	
Dairy cattle, crosses with beef breeds.....	1-57	organisms in soil.....	601-614
Damping-off diseases of conifers.....	341-348	Dormancy of seeds, relation of catalase and	
Dasheen. See <i>Colocasia esculenta</i> .		oxidase content to.....	337-341
<i>Datona integerrima</i> , effect of hydrocyanic-		Douglas fir. See <i>Pseudotsuga taxifolia</i> .	
acid gas on.....	264-268	Drought injury, compared with damping-off in	
Denitrification in diseased spinach.....	401-404	conifer seedlings.....	584-585
Desiccation, effect on catalase activity of		Drying, effect on catalase activity of seeds.....	149-150
seeds.....	149-150	Dryrot of <i>Ipomoea batatas</i>	349-350
Determining the Absolute Salt Content of		<i>Echinocystis lobata</i> , free from attacks of <i>Bac-</i>	
Soils by Means of the Freezing-Point		<i>terium lachrymans</i>	256
Method (paper).....	331-336	Effect of Carbon Disulphid and Toluol upon	
Dextrose in intumescences on frozen leaves.....	88	Nitrogen-Fixing and Nitrifying Organisms	
<i>Diablotia</i> —		(paper).....	627-634
<i>dactylis punctata</i> , possible carrier of <i>Bac-</i>		Effect of Farm Manure in Stimulating the	
<i>terium lachrymans</i>	212	Yields of Irrigated Field Crops (paper).....	402-403
<i>tritata</i> , possible carrier of <i>Bacterium lachry-</i>		Effect of Hydrocyanic-Acid Gas under	
<i>mans</i>	212	Vacuum Conditions on Subterranean	
<i>Diachasma</i> —		Larvae (paper).....	171-173
<i>fullseavi</i> —		<i>Elaeagnus</i> sp., host of damping-off parasites,	
life history.....	253-258	537-538
natural enemy of Mediterranean fruit fly		Electrometric titration of silage extract.....	125-127
.....	459-466		

Elymus—	Page		Page
<i>canadensis</i> , used in bridging-host experiments.....	236-249	Feeding stuffs, composition and energy content of.....	270
<i>macounii</i> , source of <i>Puccinia graminis tritici</i> transferred to semicongenial hosts.....	242	Fermentation, methane, in alfalfa hay and starch.....	274-275
<i>robustus</i> used in bridging-host experiment.....	234-249	<i>Feroniella lucida</i> , susceptibility to <i>Pseudomonas citri</i>	662-665
<i>virginicus</i> , used in bridging-host experiment.....	233-249	<i>Feronia limonia</i> , susceptibility to <i>Pseudomonas citri</i>	662-665
Embryo of seeds, catalase activity of.....	140-147	Feroninae, susceptibility to <i>Pseudomonas citri</i>	662-665
<i>Endophyllum semperferrini</i> , ascospore chains.....	647	Ferric oxid in—	
Endosperm of seeds, catalase activity of.....	145-147	mulch, plowsole, and subsoil.....	509-516
Energy expenditure consequent upon feed consumption.....	278-289	soil, effect of drying.....	325-328
Engelmann spruce. See <i>Picea engelmanni</i> .		Fertilizer potash residues in Hagerstown silty loam soil.....	59-81
<i>Entomophthora</i> sp., fungus attacking <i>Miris dolabratus</i>	197	Rescue, meadow. See <i>Festuca elatior</i> .	
Enzymes in alfalfa silage.....	569-590	<i>Festuca</i> —	
<i>Epithecium</i> sp., fungus isolated from <i>Ipomoea batatas</i>	357	<i>elatior</i> —	
Erb, E. S., and Frear, William (paper): Condition of Fertilizer Potash Residues in Hagerstown Silty Loam Soil.....	59-81	food plant of <i>Miris dolabratus</i>	182
<i>Eremococcus glauca</i> , susceptibility to <i>Pseudomonas citri</i>	662-665	reported bridging host for transfer of timothy rust.....	272
<i>Eriobotrya japonica</i> , host fruit of puparia parasitized by <i>Diaxasma fullawayi</i>	448	source of <i>Puccinia graminis phleipratanensis</i> transferred to semicongenial hosts.....	244
Errata and authors' emendations.....	IV	<i>pratanensis</i> , source of <i>Puccinia graminis phleipratanensis</i> transferred to semicongenial hosts.....	244
<i>Erythrpe graminis</i> , tested on <i>Bromus</i> spp.....	272	Fir, Douglas. See <i>Pseudotsuga taxifolia</i> .	
Ether extract in dry matter of feeding stuffs.....	270	Fly, fruit. See <i>Ceratitis capitata</i> .	
Ethyl-lactate method, used in determining soil acidity.....	322-329	Footroot of <i>Ipomoea batatas</i>	350
<i>Eugenia uniflora</i> —		Forests, damping-off in.....	525-558
host fruit of puparia parasitized by <i>Diaxasma fullawayi</i>	448	Formaldehyde—	
host plant of puparia parasitized by <i>Diaxasma tryoni</i>	479	effect on <i>Bacterium lactymanas</i>	205
Eugenel, used to determine oxidase activity in normal and blighted spinach.....	377-380	remedy for favus.....	417-418
Eulophinae, one species natural enemy of Mediterranean fruit fly.....	419-426	Formol, used in determining nitrogen in spinach.....	400-404
<i>Euphrasie chrysorrhoea</i> , used in experiments with <i>Limnerium validum</i>	458	<i>Fortunella</i> —	
<i>Evodia</i> —		<i>hindii</i> , susceptibility to <i>Pseudomonas citri</i>	562-605
<i>latifolia</i> , susceptibility to <i>Pseudomonas citri</i>	603-605	<i>japonica</i> , susceptibility to <i>Pseudomonas citri</i>	562-605
<i>ridleyi</i> , susceptibility to <i>Pseudomonas citri</i>	603-605	Four-lined plant bug. See <i>Poecilocapsus lineatus</i> .	
Excretion—		Frost injury to plants, growth in injected areas.....	87-88
gaseous, in alfalfa hay and starch-feeding experiment.....	273-284	Frear, William, and Erb, E. S. (paper): Condition of Fertilizer Potash Residues in Hagerstown Silty Loam Soil.....	59-81
urinary, in alfalfa hay and starch-feeding experiment.....	273-283	Freezing in plants.....	83-111
Experiments in Field Technique in Plot Tests (paper).....	251-262	Freezing-point method, determination of salt content of soil by.....	333-336
Experiments on the Value of Greensand as a Source of Potassium for Plant Culture (paper).....	483-492	Freezing points of juice from hardened and nonhardened plants.....	92-95
Fall webworm. See <i>Hyphantria cunea</i> .		French cherry. See <i>Eugenia uniflora</i> .	
Favus, skin disease caused by <i>Achorion Schönleinii</i> —		Fries, J. August, and Armsby, Henry Prentiss (paper): Net Energy Values of Alfalfa Hay and of Starch.....	269-270
anatomical changes.....	416	Frost injury—	
enzymology.....	416-417	first indications in cabbage and tomato.....	83
treatment.....	417-418	hardening cabbage against	
Feed—		chemical changes.....	95-97
effect on brisket disease.....	411-414	physiological changes.....	92-93
influence of quantity on digestibility.....	272	peroxidase changes in injured areas.....	88-90
relation to energy expenditure.....	278-280	"Frosted grains," relation to alter-ripening.....	153

Page		Page
	Fumigation with hydrocyanic acid gas, effect on subterranean larvae 133-135	Glucose in—
	Fungus parasites, cause of damping-off in conifers 511-558	hardened and nonhardened cabbage..... 95
	Further Data on the Susceptibility of Rutaceous Plants to Citrus-Canker (paper) 601-606	normal and blighted spinach..... 383
	Further Studies on Brisket Disease (paper) 499-514	Glycerin, used in determining carbohydrates in normal and blighted spinach 383
	Fusarium—	Glycamis pentaphylla, immunity to Pseudomonas citri 664-665
	<i>acuminatum</i> —	Gourd—
	fungus used in inoculation experiments with <i>Ipomoea batatas</i> 350	calabash. See <i>Lagenaria vulgaris</i> .
	not cause of damping-off in conifers..... 540-548	hedgeling. See <i>Cucumis dipsaucus</i> .
	<i>basidiola</i> , cause of damping-off in conifers..... 537-540	mandara. See <i>Cucumis acuminatus</i> .
	<i>caudatum</i> , fungus isolated from <i>Ipomoea batatas</i> 363-364	See <i>Cucurbita pepo</i> var. <i>ovifera</i> .
	<i>caluromum</i> , fungus used in inoculation experiments with <i>Ipomoea batatas</i> 359-360	Gowen, John W. (paper): Studies in Inheritance of Certain Characters of Crosses between Dairy and Beef Breeds of Cattle..... 1-78
	<i>hyperoxyperum</i> , fungus isolated from <i>Ipomoea batatas</i> 363-364	Grass—
	<i>moniliforme</i> , cause of damping-off in conifers..... 522-528	Bermuda. See <i>Cyniaria dactylon</i> .
	<i>orkhoera</i> , fungus isolated from <i>Ipomoea batatas</i> 363-364	Johnson. See <i>Sorghum halepense</i> .
	<i>oxyperum</i> —	orchard. See <i>Dactylis glomerata</i> .
	fungus isolated from <i>Ipomoea batatas</i> 360	Sudan. See <i>Andropogon halepensis</i> sudanensis.
	isolated in sweet-potato storage-rot..... 338	Tunis-sorghum hybrid, oxidase activity. 161-169
	isolated in sweet-potato storage-rot..... 360, 363-364	with. See <i>Panicum capillare</i> .
	<i>parasiticum</i> , cause of damping-off in conifers..... 537-548	Gray-damsel bug. See <i>Reduvius ferox</i>.
	<i>pinii</i> , cause of damping-off in conifers..... 537-546	Green June beetle. See <i>Cotinus nitida</i>.
	<i>solani</i> , cause of damping-off in conifers..... 540-558	Greensand, source of potassium 433-434
	<i>vaesfictum</i> —	Grub, white. See <i>Lachnosterna</i> sp.
	cause of damping-off in conifers..... 537-546	Guaiacol, used to determine oxidase activity in normal and blighted spinach 377-380
	fungus isolated from <i>Ipomoea batatas</i> 364	Guinea pig, weight of heart 431
	<i>ventricosum</i> , cause of damping-off in conifers..... 522-528	Gypsum, used in experiments with plow-sole 574-585
	Fusoma spp., reported cause of damping-off in conifers 537-546	Cypsy moth. See <i>Porticetia dispar</i> .
	Gainey, P. L. (paper): Effect of Carbon Disulphid and Tolnol upon Nitrogen-Fixing and Nitrifying Organisms 601-614	Halpin, J. G., and Beach, B. A. (paper): Observations on an Outbreak of Favus 415-418
	Galeus silvestris, parasite of Mediterranean fruit fly 431-456	Halitus citri, pest of beans 181
	Gas, hydrocyanic acid—	Hardening Process in Plants and Developments from Frost Injury (paper) 83-111
	effect on subterranean larvae..... 133-136	Hardness to frost, method of testing 93
	fumigation of cattleya orchids..... 269-268	Hardness to frost, time and temperature factors in acquiring or losing 91-94
	Gaseous excretion in alfalfa hay and starch feeding experiment 273-284	Harrington, George T., and Crocker, William (paper): Catalase and Oxidase Content of Seals in Relation to Their Dormancy, Age, Vitality, and Respiration 337-374
	Gastric digestion, effect on larvae of <i>Trikinella spiralis</i> 472-474	Harris, P. S., and Pittman, D. W. (paper): Soil Factors Affecting the Toxicity of Alkali 257-319
	Geise, Fred W., and True, Rodney H. (paper): Experiments on the value of Greensand as a Source of Potassium for Plant Culture 483-492	Harter, L. L., Weimer, J. L., and Adams, J. M. R. (paper): Sweet-Potato Storage-Rots 337-363
	Gelatin liquifiers in alfalfa silage 681-690	Hartley, Carl, and Merrill, T. C. (paper): Seedling Diseases of Conifers 521-558
	Geranium, production of injected areas due to freezing 86	Harvey, R. B. (paper): Hardening Process in Plants and Developments from Frost Injury 83-111
	Germinator, effect on catalase activity of seeds 156-159	Hawaii fruit-fly parasites in 419-426
	Germination, effect on oxidase activity of seeds 166-167	Hawkins, Lou A., et al. (paper): Physiological Studies of Normal and Blighted Spinach 369-408
	Cherkin, West Indian. <i>Cucumis anguria</i>	Hay, alfalfa—
	Gibberella saubinetii, fungus used in inoculation experiments with <i>Ipomoea batatas</i> 357-368	digestibility..... 271-282
	Glauconite, applied to farm crops 483-492	net energy value..... 269-286
	Glover, George H., and Newson, L. E. (paper): Further Studies on Brisket Disease 499-514	Hayes, H. K., and Army, A. C. (paper): Experiments in Field Tech-ic in Plot Tests 251-263

Heart—	Page	Hydroxid—Continued.	Page
condition in bricket disease.....	409-411	sodium—Continued.	
weight in—		used to—	
guinea pig.....	411	determine catalase activity of seeds.....	139
rabbit.....	411	determine water-soluble nitrogen in	
Heat production and emission in feeding		spinach.....	395-404
experiment.....	277-278, 284-286	neutralize hydrogen peroxid.....	142-144
Hedgehog gourd. See <i>Cucumis dipsaceus</i> .		Hymenoptera, found on imported orchids.....	263
<i>Helianthus annuus</i> , production of injected		<i>Hyphantria cunea</i> , host of <i>Limnerium validum</i>	458
areas due to freezing.....	86	<i>Hypochnus solani</i> . Syn. <i>Coritium vagum</i> .	
Hemiptera, found on imported orchids.....	263	<i>Hystrix patula</i> , source of <i>Puccinia graminis</i> in	
<i>Hesperethusa crenulata</i> , susceptibility to		bridging-host experiment.....	233
<i>Pseudomonas citri</i>	662-665	Inheritance of characters of crosses between	
Histological Studies on Potato Leafroll		dairy and beef cattle.....	1-57
(paper).....	559-570	Intestinal trichinae. See <i>Trichinella spiralis</i> .	
<i>Holcus halepensis sudanensis</i> . Syn. <i>Andropogon halepensis sudanensis</i> and <i>Sorghum halepensis</i> .		<i>Ipomoea batatas</i> —	
<i>Hordenium</i> —		storage-roots.....	337-368
<i>jubatum</i> —		varieties susceptible to blackrot.....	346
source of <i>Puccinia graminis tritici</i> trans-		Irish potatoes. See <i>Solanum tuberosum</i> .	
ferred to semicongenial hosts.....	249	Iron in—	
source of <i>Puccinia graminis</i> in bridging-		greensand.....	453-492
host experiment.....	230	mulch, plowsole, and subsoil.....	508-516
<i>parasilum</i> , used in bridging-host experi-		normal and blighted spinach.....	571-575
ment.....	230-249	Isoeugenol, used to determine oxidase activ-	
spp., used in bridging-host experiments.....	223-249	ity in normal and blighted spinach.....	377-386
<i>salvatore</i> —		Jack pine. See <i>Pinus banksiana</i> .	
reported infected with timothy rust		Japanese crookneck squash. See <i>Cucurbita</i>	
through bridging host.....	223	<i>moschata</i> .	
source of <i>Puccinia graminis tritici</i> trans-		Java blackrot of <i>Ipomoea batatas</i> —	
ferred to semicongenial hosts.....	247	description.....	347-348
Hubbard squash. See <i>Cucurbita maxima</i> .		inoculation experiments.....	348-349
Humic nitrogen in healthy and blighted		Jenkins, Anna E. (paper): Brown Canker of	
spinach.....	399-404	Roses, Caused by <i>Diaporthe umbrina</i>	592-600
Humus extracts of mulch, plowsole, and sub-		Jodidi, Samuel L., et al. (paper): Physiologi-	
soil, analysis.....	568	cal Studies of Normal and Blighted	
Hunter, O. W. (paper): Bacteriological		Spinach.....	369-408
Studies on Alfalfa Silage.....	571-592	Johnson grass. See <i>Sorghum halepensis</i> .	
Hydrangea, production of injected areas due		June beetle, green. See <i>Cotinus nitida</i> .	
to freezing.....	85	Kamani nuts. See <i>Terminalia catappa</i> .	
Hydrate, ammonium, used in experiments		Kellogg, Edward H., et al. (paper): Physio-	
with plowsole.....	568	logical Studies of Normal and Blighted	
Hydrochloric acid. See Acid, hydrochloric.		Spinach.....	369-408
Hydrocyanic-acid gas—		Kelly, James W., et al. (paper): Physiologi-	
effect on subterranean larvae.....	133-136	cal Studies of Normal and Blighted	
fumigation of cattleya orchids.....	263-268	Spinach.....	369-408
generation of.....	134	Kjeldahl method used in determination of	
Hydrofluoric acid. See Acid, hydrofluoric.		silage extract.....	129-131
Hydrogen—		<i>Lacknosterna</i> sp., larvae, effect of hydrocyanic-	
electrode, used in titration of silage extract.....	118-127	acid gas on.....	134-136
in gaseous excretion.....	273-284	Lactic acid. See Acid, lactic.	
peroxid, used in determining catalase		<i>Lagenaria vulgaris</i> , host plant of <i>Bacterium</i>	
content of seeds.....	140-144	<i>lachrymans</i>	206
Hydrogen-ion concentration of—		Larvae, effect of hydrocyanic-acid gas on.....	133-136
plant sap, changes in during freezing.....	98-101	<i>Lasiodiplodia nigra</i> , causal organism of rot in	
silage extract.....	218-227	<i>Ipomoea batatas</i>	347
Hydroquinone—		<i>Latua sativa</i> , production of injected areas	
oxidase of Johnson grass not active toward.....	162	due to freezing.....	86
test for oxidase in—		Lavanginae, susceptibility to <i>Pseudomonas</i>	
frozen cabbage leaves.....	88	<i>citri</i>	662-665
normal and blighted spinach.....	372-376	Leafroll of <i>Solanum tuberosum</i>	559-570
Hydroxid—		Leafspot, angular, of cucumber.....	201-220
barium, used in determining nitrogen in		Lee, H. Atherton (paper): Further data on	
spinach.....	399-404	the Susceptibility of Rutaceous Plants to	
sodium—		Citrus-Canker.....	661-666
in alfalfa silage.....	586-590	Lepidoptera, found on imported orchids.....	263

Leptolerna—
dolabrata. See *Miris dolabratus*.
id. See *Miris dolabratus*.

- Lettuce. See *Lactuca sativa*.
- Levine, M. N., et al. (paper): Plasticity of Biologic Forms of *Puccinia graminis*..... 221-249
- Levulose in intumescences on frozen leaves .. 88
- Limbe in greensand 483-492
- Limmeria validum*, natural parasite of *Hypophantria cunea*..... 428
- Loblolly pine. See *Pinus taeda*.
- Lopomorphus* id. See *Miris dolabratus*.
- Lopus* id. See *Miris dolabratus*.
- Loquat. See *Eriobotrya japonica*.
- Lycopersicon esculentum*, first indications of frost injury in..... 85
- Lygus pratensis*, general plant pest..... 181
- Macrosporium* sp., fungus isolated from *Ipomoea batatas*..... 364
- Magnesia, used in determining water-soluble nitrogen in spinach..... 368-404
- Magnesium—
in mulch, plowsole, and subsoil 368-376
oxid—
in soil, effect of drying..... 325-328
in normal and blighted spinach..... 372-375
used in determining water-soluble nitrogen in spinach..... 368-404
sulphate—
effect on plowsole..... 507
in solution used in experiments with greensand..... 486-492
- Mandarin gourd. See *Cucumis diplocarpus*.
- Manganese—
in plant tissues, relation to oxidase activity oxid—
in soil, effect of drying..... 325-328
red, in normal and blighted spinach..... 372-375
- Manganous oxid in mulch, plowsole, and subsoil..... 515-516
- Mangifera indica*, host of rot-producing organism virulent to *Ipomoea batatas*..... 347
- Mango. See *Mangifera indica*.
- Manure, effect on irrigated crops..... 493-503
- Marl, applied to farm crops..... 483-492
- Maturity of seeds, effect on—
catalase activity..... 128-140
oxidase activity..... 163-164
- Meadow fescue. See *Festuca elatior*.
- Meadow Plant Bug, *Miris dolabratus*, The (paper)..... 175-200
- Medicago sativa*, silage compared with that from *Melilotus* spp..... 113-114
- Mediterranean fruit fly. See *Ceratitis capitata*.
- Melampyrum*—
luteum, multinucleate fusion cells..... 646
pintarqua, not cause of damping-off in combers..... 550
- Melanospore* sp., fungus isolated from *Ipomoea batatas*..... 354
- Melicope triphylla*, susceptibility to *Pseudomonas citri*..... 663-665
- Melilotus* spp., silage compared with that from *Medicago sativa*..... 113-114
- Melon fly. See *Bactrocera cucurbitae*.
- Melon—
pomegranate. See *Cucumis melo* var. *dudaim*.
snake. See *Cucumis melo*.
- Mercuric-chlorid—
effect on—
Bacterium lactymanus..... 205
oxidase activity of seeds..... 168-169
used in determining nitrogen in spinach..... 399-424
nitrate, effect on oxidase activity of seeds..... 312
- Mercurous nitrate, effect on oxidase activity of seeds..... 169
- Merrill, T. C., and Hartley, Carl (paper): Seedling Diseases of Conifers..... 521-525
- Metabolism, nitrogen, in normal and blighted spinach..... 355-364
- Meta-cresol, used to determine oxidase activity in normal and blighted spinach..... 372-380
- Methane fermentation in alfalfa hay and starch..... 274-275
- Microcitrus*—
australasica, susceptibility to *Pseudomonas citri*..... 662-665
australis, susceptibility to *Pseudomonas citri*..... 662-665
- Microbial flora of alfalfa silage..... 574-592
- Mimosa elengi*, used in obtaining oviposition records for *Opus humilis*..... 443
- Miris*—
abbreviatus. See *Miris dolabratus*.
belangeri. See *Miris dolabratus*.
dolabratus—
control measures..... 197-198
description..... 191-194
dimorphism..... 183-184
distribution..... 175-181
economic importance..... 181-182
food plants..... 182-183
life history..... 184-194
natural enemies..... 194-197
synonymy..... 183
- lateralis*. See *Miris dolabratus*.
picticeps. See *Miris dolabratus*.
- Moisture—
effect on—
after-ripening of seeds..... 138
soil acidity..... 321-329
toxicity of alkali..... 366-369
in healthy and blighted spinach..... 387-390
in sweet-clover and alfalfa silage..... 114
- Molasses, effect on alfalfa silage..... 574-590
- Moneridia*—
balanina, free from attacks of *Bacterium lactymanus*..... 206
horanbia, free from attacks of *Bacterium lactymanus*..... 206
claterum, free from attacks of *Bacterium lactymanus*..... 206
- Monodactyles infusans*, causal organism of scurf in *Ipomoea batatas*..... 352
- Monocamino acid nitrogen in healthy and blighted spinach..... 399-424
- Monosaccharids in hardened and nonhardened cabbage..... 95-96
- Moss roses, resistant to brown canker..... 593-600
- Moth—
brown-tail. See *Epiproctis chrysorrhoea*.
gypsy. See *Portheia dispar*.
rusty vapor. See *Notolephus antiquus*.

	Page		Page
<i>Mucor</i> —		<i>Nolophilus antiquus</i> , used in experiments	
<i>racesanus</i> , fungus isolated from <i>Ipomoea</i>		with <i>Limnerium validum</i>	458
<i>batatas</i>	354-355	Nuts, kamanl. See <i>Terminalia catappa</i> .	
sp., not cause of damping-off in conifers.....	549	Oats. See <i>Avena sativa</i> .	
Multiple-Pipette Holder for the Distribution		Observations and Experiments on Intestinal	
of Serum for the Complement-Fixation		Trichinae (paper).....	467-474
Test, A (paper).....	615-618	Observations on an Outbreak of Favus	
<i>Murraya exotica</i> . Syn. <i>Chalcas exotica</i> .		(paper).....	415-418
Muskmelon. See <i>Cucumis melo</i> .		Olive earth, overlying greensand.....	434
<i>Nedria ipomoeae</i> , fungus isolated from		Olive, Russian wild. See <i>Elaeagnus</i> sp.	
<i>Ipomoea batatas</i>	363-364	<i>Oncophanthus binotatus</i> , infesting timothy.....	178, 181
<i>Necromyces rasilinfectum</i> , fungus isolated		Oomycetes causing damping-off of conifers.....	510
from <i>Ipomoea batatas</i>	354	Optinae, three species natural enemies of	
<i>Neofabraea malicorticis</i> , fungus used in inocu-		Mediterranean fruit fly.....	419-466
lation experiments with <i>Ipomoea batatas</i>	362	<i>Opus</i> —	
Net Energy Values of Alfalfa Hay and of		<i>felcheri</i> —	
Starch (paper).....	269-286	enemy of Mediterranean fruit fly.....	419-466
Newsom, L. E., and Glover, George H.		melen-fly parasite.....	456
(paper): Further Studies on Brisket Dis-		<i>humilis</i> —	
ease.....	409-414	life history.....	440-445
Nitrate—		natural enemy of Mediterranean fruit	
calcium, in solution used in experiments		fly.....	419-466
with greensand.....	486-492	Orchard grass. See <i>Dactylis glomerata</i> .	
mercuric, effect on oxidase activity of seeds.	169	Orchids, cattleya, fumigation with hydro-	
mercurous, effect on oxidase activity of		cyanic acid gas.....	263-268
seeds.....	166	Organic matter—	
potassium—		digestibility in alfalfa hay and starch.....	271-282
method, used in determining soil acid-		effect on toxicity of alkali.....	302-308
ity.....	322-329	Orthoptera, found on imported orchids.....	263
used in experiments with greensand.....	488-492	Osborn, Herbert (paper): The Meadow Plant	
silver, used in determining nitrogen in		Bug, <i>Miris dolabratus</i>	175-200
spinach.....	399-404	Oxid—	
sodium, effect on plowsole.....	507	aluminium, in—	
Nitric acid. See Acid, nitric.		mulch, plowsole, and subsoil.....	508-516
Nitrifying organisms in soil, effect of carbon		normal and blighted spinach.....	371-375
disulphid and toluol.....	601-614	soil, effect of drying.....	325-328
Nitrogen—		calcium, in—	
acid amid, in healthy and blighted spin-		greensand.....	383
ach.....	398-404	mulch, plowsole, and subsoil.....	515-516
amino, in alfalfa silage.....	581-590	normal and blighted spinach.....	371-375
ammonia, in healthy and blighted spin-		soil, effect of drying.....	325-328
ach.....	398-404	cuprous, in intumescences on frozen leaves.....	58
basic, in healthy and blighted spinach.....	399-404	<i>ferrie</i> , in—	
digestibility in alfalfa hay and starch.....	271-282	mulch, plowsole, and subsoil.....	508-516
humin, in healthy and blighted spinach.....	399-404	soil, effect of drying.....	325-328
in dry matter of feeding stuffs.....	270	magnesium, in—	
in hardened and nonhardened cabbage.....	97, 165-168	normal and blighted spinach.....	371-375
in silage extract.....	125-132	soil, effect of drying.....	325-328
metabolism in normal and blighted spin-		manganese, in soil, effect of drying.....	325-328
ach.....	385-404	manganous, in mulch, plowsole, and sub-	
monoamino acid, in healthy and blighted		soil.....	515-516
spinach.....	399-404	potassium, in—	
peptid, in normal and blighted spinach.....	400-404	greensand.....	485
residual soluble, in normal and blighted		normal and blighted spinach.....	371-375
spinach.....	400-404	red manganese, in normal and blighted	
Nitrogen-fixing organisms in soil, effect of		spinach.....	371-375
carbon disulphid and toluol.....	601-614	silicic, in soil, effect of drying.....	325-328
Nitrogen-free extract, digestibility in alfalfa		sodium, in normal and blighted spinach.....	371-375
hay and starch.....	270-271, 282	Oxidase, in—	
Nitrohydrochloric acid. See Acid, nitrohy-		frozen cabbage leaves.....	88
drochloric.		nonliving structures of seeds.....	167-168
Nonprotein content of dry matter of feeding		normal and blighted spinach.....	377-380
stuffs.....	270	plant tissues, relation to manganese content	
Nonprotein nitrogen in normal and blighted		seeds.....	137-174
spinach.....	393-397	Oxygen—	
Nonproteins in normal and blighted spinach.....	385-404	effect on after-ripening of seeds.....	138
Norway spruce. See <i>Picea excelsa</i> .		liberated from seeds by catalase activity.....	142-166

Page	Page
<i>Pachycrepoides dubius</i> —	Phosphate—
enemy of Mediterranean fruit fly..... 459-460	potassium, in solution used in experiments
puparium parasite of Mediterranean fruit	with greensand..... 486-492
fly..... 465-466	sodium, effect on plowsoil..... 507
<i>Panicularia pauciflora</i> , source of <i>Puccinia</i>	Phosphates in—
<i>graminis avenae</i> transferred to uncongential	greensand..... 483-492
hosts..... 243	olive earth..... 484
<i>Panicum capillare</i> , food plant of <i>Miris delab-</i>	Phosphomolybdic acid. See Acid, phospho-
<i>bratus</i> 172	molybdic.
Paracresol, used to determine—	Phosphorus—
catalase activity of seeds..... 262, 264	in hardened and nonhardened cub-
oxidase activity in normal and blighted	bage..... 97, 105-108
spinach..... 377-380	pentoxid in—
<i>Paraminyia longipedunculata</i> , susceptibility	greensand..... 485
to <i>Pseudomonas citri</i> 662-665	mulch, plowsoil, and subsoil..... 508-516
Parasites of Mediterranean fruit fly in	normal and blighted spinach..... 371-375
Hawaii..... 450-456	Phosphotungstate, barium, used in deter-
Parasitism, Morphology, and Cytology of	mining nitrogen in spinach..... 389-404
<i>Cronartium ribicola</i> (paper)..... 659-660	Phosphotungstic acid. See Acid, phospho-
Prach. See <i>Amegdalus perica</i> .	tungstic.
Pemberton, C. E., and Willard, H. F. (paper):	Physiological Studies of Normal and Blighted
Contribution to the Biology of Fruit-Fly	Spinach (paper)..... 360-408
Parasites in Hawaii..... 459-466	<i>Phytophthora</i> sp., causal organism of damping-
<i>Penicillium</i> —	off in conifers..... 530
sp., fungus isolated from <i>Ipomoea bat-</i>	<i>Pisae</i> —
<i>lar</i> 355-356, 364	<i>engelmanni</i> , host of damping-off parasites..... 531-558
sp., not cause of damping-off in conifers..... 549	<i>exola</i> , host of damping-off parasites..... 551-558
Pentoxid, phosphorus, in—	Piementel, F. J., et al. (paper): Plasticity of
greensand..... 485	Biologic Forms of <i>Puccinia graminis</i> 221-249
mulch, plowsoil, and subsoil..... 508-516	Pig, guinea, weight of heart..... 411
normal and blighted spinach..... 371-375	Pine—
<i>Pennisia officinalis</i> , host of <i>Cronartium flac-</i>	Austrian. <i>Pinus austriaca</i> .
<i>idum</i> 638-639	Corsican. See <i>Pinus nigra piceitana</i> .
Peptid nitrogen in normal and blighted	jack. See <i>Pinus banksiana</i> .
spinach..... 400-404	loblolly. See <i>Pinus taeda</i> .
<i>Peridermium</i> —	pitch. See <i>Pinus rigida</i> .
<i>brebrum</i> , parasite of <i>Pinus virginiana</i> 125	red. See <i>Pinus resinosa</i> .
<i>pini</i> —	Scotch. See <i>Pinus sylvestris</i> .
parasite on <i>Pinus sylvestris</i> 695	scrub. See <i>Pinus virginiana</i> .
<i>acrodium</i> , nuclear division of aeciospores..... 639	shortleaf. See <i>Pinus echinata</i> .
Peroxid of hydrogen, used in determining	white. See <i>Pinus strobus</i> .
catalase content of seeds..... 140-144	yellow. See <i>Pinus ponderosa</i> .
Peroxidase in frozen cabbage leaves..... 88-89	<i>Pinus</i> —
<i>Petalotia</i> —	<i>austriaca</i> , host of damping-off parasites..... 531-558
<i>funerea</i> , isolated from damped-off conifers..... 547-558	<i>banksiana</i> , host of damping-off parasites..... 528-558
<i>haripii</i> , isolated from damped-off conifers..... 547-558	<i>echinata</i> , germination in mineral soil..... 526
sp., isolated from <i>Ipomoea batatas</i> 264	<i>flexilis</i> , host of <i>Cronartium ribicola</i> 620
<i>Phaeole neogaeophala</i> , natural enemy of Med-	<i>montana</i> , attacked by <i>Melampsora pin-</i>
iterranean fruit fly..... 459-460	<i>torqua</i> 551
Phenolphthalein, used in—	<i>nigra piceitana</i> , host of damping-off
determining catalase content of seeds..... 339	fungi..... 524-558
titration of silage extract..... 216-217	<i>parviflora</i> —
<i>Phicum pratense</i> —	host of <i>Cronartium ribicola</i> 630
food plant of <i>Miris delabratus</i> 276, 182-183	nature of infection by <i>Cronartium ribi-</i>
source of <i>Puccinia graminis phaeipratis</i>	<i>cola</i> 649, 651
transferred to semicongenial hosts..... 244	<i>pinus-trispora</i> —
uncongential host of <i>Puccinia graminis</i>	host of damping-off parasites..... 521-558
<i>avenae</i> 243	nature of injury from <i>Cronartium comp-</i>
Phloroglucin, used to determine oxidase	<i>tenosa</i> 649
reaction in normal and blighted spinach..... 372-380	<i>resinosa</i> , host of damping-off parasites..... 531-558
<i>Phoma</i> —	<i>resida</i> , germination in mineral soil..... 526
<i>batatas</i> , reported causal organism of dryrot	<i>scopulorum</i> . Syn. <i>Pinus ponderosa</i> .
of <i>Ipomoea batatas</i> 549	<i>strobus</i> —
<i>betae</i> , not cause of damping-off in conifers..... 550-558	host of <i>Cronartium ribicola</i> 610-660
<i>Phoranthia occidentis</i> , parasite of <i>Miris del-</i>	host of damping-off parasites..... 530-558
<i>bratus</i> 170	<i>lyceitris</i> , host of—
	damping-off parasites..... 531-558
	<i>Peridermium pini</i> 625

Pinus—Continued.	Page	Page
<i>taeda</i> , germination in mineral soil.....	506	Potassium-alum solution, used in determining protein nitrogen in spinach..... 391-393
<i>virginiana</i> —		Potassium-nitrate method, used in determining soil acidity..... 322-339
germination in mineral soil.....	506	Potato—
host of <i>Peridermium cerebrum</i>	625	sweet. See <i>Ipomoea batatas</i> .
Pipette holder for distribution of serum.....	615-618	See <i>Solanum tuberosum</i> .
Pitch pine. See <i>Pinus rigida</i> .		Potentiometric determination of changes in hydrogen-ion concentration of plant sap during freezing..... 99-101
Pittman, D. W., and Harris, F. S. (paper):		Protein—
Soil Factors Affecting the Toxicity of Alkali.....	287-319	digestibility in alfalfa hay and starch..... 271, 287
Plant bug—		in dry matter of feeding stuffs..... 270
four-lined. See <i>Poecilocarpus lineatus</i> .		in hardened and nonhardened cabbage. 97, 103-104
meadow. See <i>Miris dolabratus</i> .		in normal and blighted spinach..... 385-404
tarnished. See <i>Lygus pratensis</i> .		<i>Pseudomonas</i> —
Plants, hardening against frost.....	83-111	<i>citri</i> —
Plasmolysis of cells—		susceptibility of rutaceous plants..... 661-666
caused by freezing.....	90-92	virulence to Rutaceae..... 661-666
effect on hydrogen-ion concentration of sap.....	99	<i>lachrymans</i> . Syn. <i>Bacterium lachrymans</i> .
Plasticity of Biologic Forms of Puccinia graminis (paper).....	221-249	Puccinia—
<i>Plenodomus destruens</i> , causal organism of foot-rot of <i>Ipomoea batatas</i>	350	<i>dispersa</i> , plasticity of biologic forms..... 221
Plot tests, field technic.....	251-263	<i>falsaria</i> , aeciospore chains..... 647
Flowrate, relation of inorganic soil colloids to.....	593-519	<i>graminis</i> —
<i>Poa pratensis</i> —		plasticity of biologic forms..... 221-249
catalase and oxidase content of seeds.....	137-174	<i>agrostis</i> , plasticity of biologic forms..... 241-247
food plant of <i>Miris dolabratus</i>	183	<i>avenae</i> , used in bridging-host experiments..... 227-249
<i>Poecilocarpus lineatus</i> , pest of currants.....	181	<i>phlepratensis</i> , plasticity of biologic forms..... 241-247
Polypeptids in normal and blighted spinach.....	385-404	<i>secalis</i> , used in bridging-host experiments..... 225
<i>Polyphagus euglenae</i> , similarity to <i>Cronarium ribicola</i>	648	<i>tritici</i> , used in bridging-host experiments..... 225-249
Polysaccharids in hardened and nonhardened cabbage.....	95-96	<i>tritici-compacti</i> , used in bridging-host experiments..... 227-249
Pomegranate melon. See <i>Cucumis melo</i> var. <i>dudaim</i> .		<i>Pucciniastrum agrimoniae</i> , structure of ure-dinium..... 662
<i>Popillia japonica</i> , larvae, effect of hydrocyanic acid gas on.....	134-146	Pulverization of seeds, effect on catalase activity..... 143-145
<i>Populus</i> —		Pumpkin. See <i>Cucurbita pepo</i> .
<i>alba</i> , attacked by <i>Melampora pinioregna</i>	551	Pyrocatechin, used as test for oxidase in frozen cabbage leaves..... 88
<i>balsamifera candicans</i> , attacked by <i>Melampora pinioregna</i>	551	Pyrocatechol, used to determine oxidase activity of—
<i>tremula</i> , attacked by <i>Melampora pinioregna</i>	551	normal and blighted spinach..... 377-380
<i>Porteria dispar</i> , used in experiments with <i>Limnerium radium</i>	458	seeds..... 162, 166
Potash in—		Pyrogallol, used to determine oxidase in—
greensand.....	483-492	frozen cabbage leaves..... 88
Hagerstown silty loam, soluble in—		normal and blighted spinach..... 377-380
ammonium-chlorid solution.....	67-70	seeds..... 162, 166
carbonated water.....	67	<i>Pseudotsuga latifolia</i> , host of damping-off parasites..... 531-558
distilled water.....	64-66	<i>Pythium debaryanum</i> , cause of damping-off in conifers..... 524-558
hydrochloric acid.....	63-64	Rabbit, weight of heart..... 411
Potassium—		"Rain-barley," relation to after-ripening..... 153
chlorid, used in experiments with green-sand.....	488	Rambler roses, resistant to brown canker..... 553-560
cyanid, tolerance of larvae of <i>Trichinella spiralis</i> to.....	477-479	Reagents used in determining oxidase activity of seeds..... 162, 166
in greensand.....	483-492	Red clover. See <i>Trifolium pratense</i> .
nitrate, used in experiments with green-sand.....	488-492	Red manganese oxid in normal and blighted spinach..... 374-375
oxid in—		Red pine. See <i>Pinus resinosa</i> .
normal and blighted spinach.....	371-375	Reducing sugars in—
greensand.....	485	hardened and nonhardened cabbage..... 95-96
phosphate in solution used in experiments with greensand.....	486-492	healthy and blighted spinach..... 381-384
		<i>Redusolusferus</i> , natural enemy of <i>Miris dolabratus</i> 194, 196

	Page		Page
Relation of Inorganic Soil Colloids to Plow- sole in Citrus Groves in Southern Cali- fornia.....	505-519	<i>Secale cereale</i> , used in bridging-host experi- ments.....	246-249
Residual soluble nitrogen in normal and blighted spinach.....	400-404	Seed, dormant, decay due to damping-off.....	525
Respiration of seeds, relation of catalase and oxidase content to.....	537-574	Seedling Diseases of Conifers (paper).....	521-558
Reynolds, Francois H. (paper): A Multiple- Pipette Holder for the Distribution of Serum for the Complement-Fixation Test.....	615-618	Seedlings, conifer, injury from— damping-off.....	521-558
<i>Rhectoparacanium aphanidermatum</i> , causal organism of damping-off in conifers.....	530-540	drouth.....	554-555
Rhizoctonia, causal organisms of damping-off in conifers.....	522-538	mechanical means.....	551-554
<i>Rhizoctonia</i> — <i>crocorum</i> , not grown on artificial media....	530	whitespot.....	551
<i>mediaginis</i> , Syn., <i>Rhizoctonia crocorum</i> .		Seeds— catalase and oxidase content of.....	537-574
<i>Rhizopus nigricans</i> , causal organism of sweet- potato storage-rots.....	338-368	effect of— after-ripening on catalase activity.....	152-156
<i>Ribes</i> spp., hosts of <i>Cronartium ribicola</i>	619-660	age on catalase activity.....	155-166
Ringrot of <i>Ipomoea batatas</i> — dissemination.....	347	age on oxidase activity.....	150-154
Inoculation experiments.....	347-344	brometin on oxidase activity.....	164-165
symptoms.....	340-343	drying on catalase activity.....	160
<i>Rosa</i> spp., resistant to brown canker.....	593-600	germination on oxidase activity.....	149-150
<i>Comodysium fuchetii</i> , difference from <i>Diophor-</i> <i>the umbrina</i>	591-594	heat on catalase activity.....	166-167
<i>Rosellinia</i> sp., not cause of damping-off in conifers.....	556	maturity on catalase activity.....	154-156
Roses. See <i>Rosa</i> spp.		maturity on oxidase activity.....	163-164
Rugosa, resistant to brown canker.....	593-600	mercuric chlorid on oxidase activity.....	168-166
Russian wild olive. See <i>Elaeagnus</i> sp.		mercuric nitrate on oxidase activity.....	169
Rust, timothy, transferred by bridging hosts.....	577	mercurous nitrate on oxidase activity.....	169
Rusty vaporermoth. See <i>Notodiphus anticus</i> .		retention in germinator on catalase activ- ity.....	150-159
Rutaceous plants, susceptibility to citrus canker.....	661-666	vitality on catalase activity.....	150-154
•ye— effect on alfalfa silage.....	574-590	oxidase activity of nonliving structures.....	165-168
See <i>Secale cereale</i> .		Serum, distributed by multiple-pipette holder.....	615-618
Salt content of soils, determination by freezing-point method.....	531-536	<i>Severinia busifolia</i> , immunity to <i>Pseud-</i> <i>monas citri</i>	562-605
Salvia, production of injected areas due to freezing.....	86	Shortleaf pine. See <i>Pinus echinata</i> .	
Sanford, H. L., and Sasser, E. R. (paper): Effect of Hydrocyanic-Acid Gas under Vacuum Conditions on Subterranean Larvæ.....	133-156	Silage— alfalfa— acidity.....	574-584
Sasser, E. R., and Dietz, H. F. (paper): Fumigation of Cattleya Orchids with Hy- drocyanic-Acid Gas.....	263-268	bacteria in.....	571-592
Sasser, E. R., and Sanford, H. L. (paper): Effect of Hydrocyanic-Acid Gas under Vacuum Conditions on Subterranean Larvæ.....	133-156	bacteriological analysis.....	586-587
Schwartz, Benjamin (paper): Observations and Experiments on Intestinal Trichinæ.....	467-482	chemical analysis.....	580-587
<i>Sclerotium</i> — <i>bataticola</i> , causal organism of charcoal-rot in <i>Ipomoea batatas</i>	351	effect of carbohydrates.....	584-590
<i>rolfsii</i> , fungus isolated from <i>Ipomoea batatas</i> .	364	loss in weight.....	114
Scotfield, C. S. (paper): Effect of Farm Manure in Stimulating the Yields of Irrig- ated Farm Crops.....	491-503	moisture content.....	114
Scotch pine. See <i>Pinus sylvestris</i> .		sweet-clover compared with alfalfa.....	113-115
Scrub pine. See <i>Pinus virginiana</i> .		Silica in— greensand.....	485-495
Scurf, fungous disease of <i>Ipomoea batatas</i>	352-353	mulch, plowsole, and subsoil.....	508-516
		Silico-oxid content of soil, effect of drying.....	525-528
		Silicon dioxide in— mulch, plowsole, and subsoil.....	508-516
		normal and blighted spinach.....	571-575
		Silver— chlorid, used in determining nitrogen in spinach.....	429-434
		nitrate, used in determining nitrogen in spinach.....	399-404
		Snake melon. See <i>Cucumis melo</i> .	
		Sodium— carbonate— effect on plowsole.....	597
		used in alkali experiment.....	599-579
		chlorid— effect on plowsole.....	597
		used in alkali experiment.....	599-579
		used in experiments with intestinal trichinæ.....	468-482

Sodium—Continued.	Page	Stakman, E. C., Piemeisel, F. J., and Levine, M. N. (paper): Plasticity of Biologic Forms of <i>Puccinia graminis</i>	Page
cyanid—		221-249
effect on cattleya orchids.....	267-268	Starch—	
used to generate hydrocyanic acid gas.....	134	digestibility.....	271, 282
hydroxid—		influence on digestibility of hay.....	272
in alfalfa silage.....	556-590	in hardened and nonhardened cabbage.....	95-96, 105-108
used to determine—		in healthy and blighted spinach.....	381-384
catalase content of seeds.....	139	methane fermentation.....	274-275
water-soluble nitrogen in spinach.....	398-404	net energy value.....	269-286
nitrate, effect on plowsole.....	597	Storage, effect on oxidase activity of seeds.....	165-166
oxid in normal and blighted spinach.....	371-375	Storage-rots of <i>Ipomoea batatas</i>	337-368
phosphate, effect on plowsole.....	597	Straw, effect on alfalfa silage.....	574-590
sulphate, used in alkali experiment.....	289-319	Studies in Inheritance of Certain Characters of Crosses between Dairy and Beef Breeds of Cattle (paper).....	1-57
Softrot in <i>Ipomoea batatas</i> —		Stutzer's nitrogen in silage.....	129-132
dissemination.....	142	Sucrose in—	
inoculation experiments.....	142-144	hardened and nonhardened cabbage.....	95-96
symptoms.....	140-142	healthy and blighted spinach.....	381-384
Soil Acidity as Affected by Moisture Conditions of the Soil (paper).....	322-329	Sudan grass. See <i>Andropogon halepensis sudanensis</i> .	
Soil—		Sugar beets. See <i>Beta vulgaris</i> .	
adobe, inorganic colloids.....	516	Sugar, cane, effect on alfalfa silage.....	584-590
colloids, relation to plowsole.....	595-519	Sugars in healthy and blighted spinach.....	381-384
effect of carbon disulphid and toluol on nitrogen-fixing organisms.....	601-614	Sulphate—	
Hagerstown silty loam, fertilizer potash residues in.....	59-81	calcium, effect on plowsole.....	597
overwintering of <i>Bacterium lactymani</i> in.....	213	copper, effect on <i>Bacterium lactymani</i>	205
particles, effect of size on toxicity of alkali.....	290-301	magnesium—	
salt content of, determination by freezing-point method.....	331-336	effect on plowsole.....	597
water content of, effect on fumigation with hydrocyanic acid gas.....	133-136	in solution used in experiments with greensand.....	486-492
Soil Factors Affecting the Toxicity of Alkali (paper).....	287-319	sodium, used in alkali experiment.....	289-290
<i>Solanum tuberosum</i> —		Sulphid, copper, used in determining nitrogen in spinach.....	400-402
effect of farm manure on.....	494-593	Sulphur trioxid in—	
leafroll.....	559-570	greensand.....	485
<i>Sorghum halepensis</i> —		normal and blighted spinach.....	371-375
diameter of cells in seeds.....	145	Sulphuric acid. See Acid, sulphuric.	
catalase and oxidase content of seeds.....	137-174	Sunflower. See <i>Helianthus annuus</i> .	
hybridized with Tunis grass, catalase activity of seeds.....	146-147	Swanson, C. O., and Tague, E. L. (paper): Chemistry of Sweet-Clover Silage in Comparison with Alfalfa Silage.....	113-132
Species, new.....	593-600	Sweet clover. See <i>Medicago</i> spp.	
<i>Sphaeromena fimbriatum</i> , causal organism of blackrot of <i>Ipomoea batatas</i>	344	Sweet-clover silage compared with alfalfa silage.....	113-132
<i>Spinacia oleracea</i> —		Sweet-Potato Storage-Rots (paper).....	337-368
denitrification due to blight.....	401-404	Tague, E. L., and Swanson, C. O. (paper): Chemistry of Sweet-Clover Silage in Comparison with Alfalfa Silage.....	113-132
effect of blight on—		Tarnished plant bug. See <i>Lycus pratensis</i> .	
ash content.....	371-375	Tarax—	
carbohydrate production.....	381-384	<i>canadensis</i> , host of damping-off parasites.....	531-558
nitrogen metabolism.....	385-404	<i>cuspidata</i> , host of damping-off parasites.....	531-558
oxidase reaction.....	377-380	Temperature—	
Spinach-blight, physiological studies.....	369-408	effect on—	
<i>Sporobolus cryptandrus</i> , source of <i>Puccinia graminis</i> in bridging-host experiment.....	2-6	<i>Bacterium lactymani</i>	204-205
<i>Sporotrichum</i> sp., fungus isolated from <i>Ipomoea batatas</i>	384	catalase activity of seeds.....	154-156
Spurge—		relation to hardening of plants against frost injury.....	91-94
Engelmann. See <i>Picea engelmannii</i> .		<i>Terminalia catappa</i> , host of puparia parasitized by <i>Diachasma tryoni</i>	449
Norway. See <i>Picea excelsa</i> .		Tetramethylparaphenylenediamin, used as a test for oxidase in frozen cabbage leaves.....	88
Squash—			
Hubbard. See <i>Cucurbita maxima</i> .			
Japanese crookneck. See <i>Cucurbita moschata</i> .			
summer. See <i>Cucurbita pepo</i> var. <i>condensa</i> .			
Stirring cucumber. See <i>Momordica charitri-</i>			

	Page		Page
<i>Tetrastichus giffardianus</i> —		True, Rodney H., Black, Otis F., Kelly, James W., Bunzell, H. H., Hawkins, Lon A., Jodidi, Samuel L., and Kellogg, Edward H. (paper): Physiological Studies of Normal and Blighted Spinach.....	369-378
life history.....	448-454	True, Rodney H., and Geise, Fred W. (paper): Experiments on the Value of Greensand as a Source of Potassium for Plant Culture.....	483-492
natural enemy of Mediterranean fruit fly.....	419-426	Tunis grass and sorghum hybrid, catalase activity of seeds.....	149-147
<i>Thevetia nerifolia</i> —		Turban squash. <i>Cucurbita pepo</i> var. <i>condensata</i>	
host of puparia parasitized by <i>Diachasma fullawayi</i>	448	Tyrosin—	
host of larvae parasitized by <i>Tetrastichus giffardianus</i>	452	used to determine oxidase activity in normal and blighted spinach.....	377-380
Thymolphthalein, used in titration of silage extract.....	119-127	test for oxidase in frozen cabbage leaves.....	88
<i>Tilia americana</i> , catalase and oxidase content of seeds.....	138	Tyrosine, oxidase of Johnson grass not active toward.....	162
Timothy—		Urinary excretion, analysis in alfalfa hay and starch-feeding experiment.....	273-283
rust, transferred by bridging hosts.....	222	Vacuum process of fumigation with hydrocyanic acid gas.....	133-136
See <i>Phleum pratense</i>		Vaseline, used in treatment of larvae.....	417-418
Titration of silage extract.....	110-130	<i>Verticillium cinnabarinum</i> , fungus isolated from <i>Ipomoea batatas</i>	364
<i>Toddalia asiatica</i> , susceptibility to <i>Pseudomonas citri</i>	663-665	Vitality of seeds, relation of catalase and oxidase content to.....	137-174
Toluol—		Water in soil, effect on fumigation with hydrocyanic acid gas.....	133-136
effect on nitrogen-fixing organisms in soil.....	601-614	Water extract of silage—	
used in determining carbohydrates in normal and blighted spinach.....	383	acidity.....	116-125
Tomato. See <i>Lycopersicon esculentum</i>		method of making.....	114-115
Toxicity of alkali, effect of soil factors.....	287-310	nitrogen in.....	125-132
<i>Trichinella spiralis</i> , growth and molting.....	467-483	Watermelon. See <i>Citrullus vulgaris</i>	
<i>Trichoderma</i> —		Wax on plant leaves, effect upon freezing point of plant.....	86
<i>koningi</i> , isolated from—		Webworm, fall. See <i>Hyphantria cunea</i>	
damped-off conifers.....	546-558	Weimer, J. L., et al. (paper): Sweet-Potato Storage-Rots.....	337-368
<i>Ipomoea batatas</i>	360-361	West Indian gherkin. See <i>Cucumis anguria</i>	
isolated from damped-off conifers.....	519-558	Wheat. See <i>Triticum aestivum</i>	
used in inoculation experiments with <i>Ipomoea batatas</i>	361	Wheat wireworm. See <i>Agriotes mancus</i>	
<i>Trichosanthes colubrina</i> , free from attacks of <i>Bacterium lactyrans</i>	206	White grub. See <i>Larimorpha</i> sp.	
<i>Trichosporium</i> sp., fungus isolated from <i>Ipomoea batatas</i>	361	White pine. See <i>Pinus strobus</i>	
<i>Trichothecium roseum</i> , not cause of damping-off in conifers.....	540-558	Whitespot, resemblance to damping-off in conifer seedlings.....	551
<i>Trifolium</i> —		Wild cucumber. See <i>Echinocystis lobata</i>	
<i>incarnatum</i> —		Willard, H. F., and Pemberton, C. E. (paper): Contribution to the Biology of Fruit-Fly Parasites in Hawaii.....	419-466
catalase content of seeds.....	144	Wireworm, wheat. See <i>Agriotes mancus</i>	
diameter of cells in seeds.....	145	Witch grass. See <i>Panicum capillare</i>	
<i>pratense</i> , used in experiments with greensand.....	488-494	<i>Xanthoxylum rheticum</i> , immunity to <i>Pseudomonas citri</i>	663-665
Trioxid, sulphur, in—		Yeasts in alfalfa silage.....	574-592
greensand.....	488	Yellow pine. See <i>Pinus ponderosa</i>	
normal and blighted spinach.....	377-378	Yew. See <i>Taxus cuspidata</i>	
<i>Trifolium trifolium</i> , immunity to <i>Pseudomonas citri</i>	662-665	<i>Zygothynchus</i> sp., fungus isolated from <i>Ipomoea batatas</i>	364
<i>Tritium</i> —		Zymase, cause of alcohol in silage.....	590
<i>aestivum</i> —			
catalase activity of parts of seed.....	140		
used in bridging-host experiments.....	226-249		
used in experiments with greensand.....	488-492		
yield on alkali land.....	302		
<i>vulgare</i> , reported infected with timothy rust through bridging host.....	222		



